

**Exercise sheet no. 9 – Numerics for instationary differential equations**

**Exercise 23:**

Solve the one-dimensional wave equation  $u_{tt} = c^2 u_{xx}$  on  $[0, \pi]$  with initial values

$$u(x, 0) = u_0(x), \quad u_t(x, 0) = v_0(x)$$

and homogeneous Neumann boundary conditions

$$u_x(0, t) = u_x(\pi, t) = 0$$

using Fourier series. For this, assume that the exact solution  $u$  exists and extend it symmetrically to  $[-\pi, 0]$ . Prove: if  $u_0$  and  $v_0$  are real-valued, then  $u$  is real-valued.

**Exercise 24:**

Consider the differential equation  $u_t = cu_x$  and the Lax-Friedrichs method

$$\frac{u_j^{n+1} - \frac{1}{2}[u_{j+1}^n + u_{j-1}^n]}{\tau} = c \frac{u_{j+1}^n - u_{j-1}^n}{2h}$$

as well as the Lax-Wendroff method

$$\frac{u_j^{n+1} - u_j^n}{\tau} = c \frac{u_{j+1}^n - u_{j-1}^n}{2h} + \frac{c^2 \tau}{2} \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{h^2}.$$

For the initial value  $u(x, 0) = \exp(i\alpha x)$ , determine the growth factor  $G(\alpha)$  and do a von Neumann stability analysis, that is, formulate a condition for  $c\tau/h$ , such that  $|G(\alpha)| \leq 1$  for all  $\alpha \in \mathbb{R}$ .

**Programming exercise 2 :**

Implement the two methods of the previous exercise for the one-dimensional problem

$$\begin{aligned} u_t(x, t) &= cu_x(x, t), & x \in [x_{\min}, x_{\max}], t > 0, \\ u(x, 0) &= \alpha \exp(-\beta(x - \gamma)^2), & x \in [x_{\min}, x_{\max}]. \end{aligned}$$

- Use as boundary conditions

$$u(x_{\min}, t) = \alpha \exp(-\beta(x_{\min} + ct - \gamma)^2), \quad u(x_{\max}, t) = \alpha \exp(-\beta(x_{\max} + ct - \gamma)^2).$$

- Experiment with different values of  $\alpha, \beta$  and  $\gamma$  and make clear how these parameters affect the solutions. Write a short comment.
- Test different values of  $c$ . In which direction the wave is transported?
- Experiment with different values of  $c, \tau$  and  $h$  to see whether the numerical solution is bounded or not. How does it coincide with the von Neumann stability analysis of exercise 25?
- Create convergence plots for  $h$ , for instance using  $c = -0.5, \tau = \frac{1}{160}$  and

$$h = \frac{1}{10}, \frac{1}{20}, \frac{1}{40}, \frac{1}{80}.$$

Integrate until  $N\tau = t_{\text{end}} = 0.5$  and plot the errors  $\max_j |u(x_j, t_{\text{end}}) - u_j^N|$ . As parameters, choose

$$\alpha = 1, \quad \beta = 10, \quad \gamma = 0.2, \quad x_{\min} = 0, \quad x_{\max} = 1.$$

**Solutions are discussed on 10.07.2024.**

**The programming exercise needs to be handed in by July 17th, noon.**

Contact person: Dominik Sulz - [dominik.sulz@uni-tuebingen.de](mailto:dominik.sulz@uni-tuebingen.de). Open door policy - just come to my office if you have any questions!