

CONVERGENCE OF A FLUX-SPLITTING FINITE VOLUME SCHEME FOR CONSERVATION LAWS DRIVEN BY LÉVY NOISE.

ANANTA K. MAJEE

ABSTRACT. We explore numerical approximation of multidimensional stochastic balance laws driven by multiplicative Lévy noise via flux-splitting finite volume method, and establish the convergence of such approximations to an entropy solution for the underlying problem.

1. INTRODUCTION

Let $(\Omega, \mathbb{P}, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0})$ be a filtered probability space satisfying the usual hypothesis i.e $\{\mathcal{F}_t\}_{t \geq 0}$ is a right-continuous filtration such that $\mathcal{F}_0$ contains all the $\mathbb{P}$ -null subsets of $(\Omega, \mathcal{F})$. In this paper, we are interested in the study of numerical scheme and numerical approximation for the multi-dimensional nonlinear stochastic balance laws of type

$$\begin{aligned} du(t, x) + \operatorname{div}_x(\vec{v}(t, x)f(u(t, x))) dt &= \int_{\mathbf{E}} \eta(u(t, x); z) \tilde{N}(dz, dt), \quad (t, x) \in \Pi_T, \\ u(0, x) &= u_0(x), \quad x \in \mathbb{R}^d, \end{aligned} \quad (1.1)$$

where $\Pi_T = [0, T) \times \mathbb{R}^d$ with $T > 0$ fixed. Here, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a given real valued flux function, $\vec{v}$ is a given vector valued function, $u_0(x)$ is given initial function and $\tilde{N}(dz, dt) = N(dz, dt) - m(dz) dt$, where N is a Poisson random measure on $(\mathbf{E}, \mathcal{E})$ with intensity measure $m(dz)$, where $(\mathbf{E}, \mathcal{E}, m)$ is a σ -finite measure space. Furthermore, $(u, z) \mapsto \eta(u, z)$ is a given real valued functions signifying the multiplicative nature of the noise.

This type of equation arises in many different fields where non-Gaussianity plays an important role. As for example, it has been used in models of neuronal activity accounting for synaptic transmissions occurring randomly in time as well as at different locations on a spatially extended neuron, chemicals reaction-diffusion systems, market fluctuations both for risk management and option pricing purpose, stochastic turbulence, etc. The study of well-posedness theory for this kind of equation is of great importance in the light of current applications in continuum physics.

Remark 1.1. We will carry out our analysis under the structural assumption $\mathbf{E} = \mathcal{O} \times \mathbb{R}^*$ where $\mathcal{O}$ is a subset of the Euclidean space. The measure m on $\mathbf{E}$ is defined as $\lambda \times \mu$ where λ is a Radon measure on $\mathcal{O}$ and μ is so-called Lévy measure on $\mathbb{R}^*$. Such a noise would be called an impulsive white noise with jump position intensity λ and jump size intensity μ . We refer to [29] for more on Lévy sheet and related impulsive white noise.

In the case $\eta = 0$, the equation (1.1) becomes a standard conservation laws in $\mathbb{R}^d$ and there exists a satisfactory well-posedness theory based on Kruzkov's pioneering idea to picks up the physically relevant solution in an unique way, called *entropy solution*. We refer to [19, 26, 27, 31] and references therein for more on entropy solution theory for deterministic conservation laws.

The study of stochastic balance laws driven by noise is comparatively new area of pursuit. Only recently balance laws with stochastic forcing have attracted the attention of many authors [2, 4, 5, 8, 9, 10, 11, 12, 14, 15, 16, 18, 21, 22] and resulted in significant momentum in the theoretical development of such problems. Due to nonlinear nature of the underlying problem, explicit solution formula is hard to obtain and hence robust numerical schemes for approximating such equation are very important. In the

2000 *Mathematics Subject Classification.* 45K05, 46S50, 49L20, 49L25, 91A23, 93E20.

Key words and phrases. Conservation Laws, Stochastic Forcing, Lévy Noise, Stochastic Entropic Solution, Stochastic Partial Differential Equations, Young measures, Kruzkov's entropy, Finite volume scheme.

last decade, there has been a growing interest in numerical approximation and numerical experiments for entropy solution to the related Cauchy problem driven by stochastic forcing. The first documented development in this direction is [20], where the authors established existence of weak solution (possibly non-unique) of one dimensional balance law driven by Brownian noise via splitting method. In a recent paper [25], Kröker and Rodhe established the convergence of monotone semi-discrete finite volume scheme by using stochastic compensated compactness method. Buazet [3] revisited the paper of Holden and Risebro [20], and generalized the operator-splitting method for the same Cauchy problem but in a bounded domain of $\mathbb{R}^d$. Using Young measure theory, author established the convergence of approximate solutions to an entropy solution. We also refer to see [24], where the time splitting method was analyzed for more general noise coefficient in the spirit of Malliavin calculus and Young measure theory. In a recent papers [6, 7], Bauzet et. al. have studied fully discrete scheme via flux-splitting and monotone finite volume schemes for stochastic conservation law driven by multiplicative Brownian noise and established its convergence by using Young measure technique.

The study of numerical schemes for stochastic balance laws driven by Lévy noise is more sparse than the previous case. A semi-discrete finite difference scheme for a conservation laws driven by a homogeneous multiplicative Lévy noise has been studied by Koley et al.[23]. Using BV estimates, the authors showed the convergence of approximate solutions, generated by the finite difference scheme, to the unique entropy solution as the spatial mesh size $\Delta x \rightarrow 0$ and established rate of convergence which is of order $\frac{1}{2}$.

The above discussions clearly highlights the lack of the study of fully discrete scheme and its convergence for stochastic balance laws driven by Lévy noise. In this paper, drawing primary motivation from [6], we propose a fully discrete flux-splitting finite volume scheme for (1.1), and address the convergence of the scheme. First we establish few essential *a-priori* estimates for approximate solutions and then using these estimates, we deduce entropy inequality for approximate solutions. Using Young measure theory, we conclude that the finite volume approximate solutions tend to a generalized entropy solution of (1.1).

The rest of the paper is organized as follows. In Sections 2 and 3, we collect all the assumptions for the subsequent analysis, define proposed numerical scheme and finally state the main result of this article. Section 4 deals with few a priori estimates on the finite volume approximate solutions and using these a priori estimates, in Section 5, we establish discrete and continuous version of entropy inequalities on approximate solutions. The Section 6 is devoted for the proof of the main theorem along with short discussion of Young measure theory and its compactness, is presented in Appendix 7.

2. PRELIMINARIES AND TECHNICAL FRAMEWORK

It is well-known that due to nonlinear flux term in (1.1), solutions to (1.1) are not necessarily smooth even if initial data is smooth, and hence must be interpreted via weak sense. Before introducing the concept of weak solutions, we first assume that $(\Omega, P, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0})$ be a filtered probability space satisfying the usual hypothesis, i.e., $\{\mathcal{F}_t\}_{t \geq 0}$ is a right-continuous filtration such that $\mathcal{F}_0$ contains all the P -null subsets of $(\Omega, \mathcal{F})$. Moreover, by a predictable σ -field on $[0, T] \times \Omega$, denoted by $\mathcal{P}_T$, we mean that the σ -field generated by the sets of the form: $\{0\} \times A$ and $(s, t] \times B$ for any $A \in \mathcal{F}_0; B \in \mathcal{F}_s$, $0 < s, t \leq T$. The notion of stochastic weak solution is defined as follows:

Definition 2.1 (Weak solution). An $L^2(\mathbb{R}^d)$ -valued $\{\mathcal{F}_t : t \geq 0\}$ -predictable stochastic process $u(t) = u(t, x)$ is called a stochastic weak solution of (1.1) if for all nonnegative test functions $\psi \in C_c^\infty([0, T] \times \mathbb{R}^d)$,

$$\begin{aligned} \int_{\mathbb{R}^d} \psi(0, x) u_0(x) dx + \int_{\Pi_T} \left\{ \partial_t \psi(t, x) u(t, x) + \vec{v}(t, x) f(u(t, x)) \cdot \nabla_x \psi(t, x) \right\} dt dx \\ + \int_{\Pi_T} \int_{\mathbf{E}} \eta(u(t, x); z) \psi(t, x) \tilde{N}(dz, dt) dx = 0, \quad \mathbb{P} - \text{a.s.} \end{aligned}$$

However, since there are infinitely many weak solutions, one needs to define an extra admissibility criteria to select physically relevant solution in a unique way, and one such condition is called entropy condition. Let us begin with the notion of entropy flux pair.

Definition 2.2 (Entropy flux pair). (β, ϕ) is called an entropy flux pair if $\beta \in C^2(\mathbb{R})$ and $\zeta : \mathbb{R} \mapsto \mathbb{R}$ such that

$$\zeta'(r) = \beta'(r) f'(r).$$

An entropy flux pair (β, ζ) is called convex if $\beta''(s) \geq 0$.

Let $\mathcal{A} = \{\beta \in C^2(\mathbb{R}), \text{convex such that support of } \beta'' \text{ is compact}\}$. In the sequel, we will use specific entropy flux pairs. For any $a \in \mathbb{R}$ and $\beta \in \mathcal{A}$, define $F^\beta(a) = \int_0^a \beta'(s) f'(s) ds$. Note that, $F^\beta(\cdot)$ is a Lipschitz continuous function on $\mathbb{R}$ and (β, F^β) is an entropy flux-pair. To this end, we define the notion of stochastic entropy solution of (1.1).

Definition 2.3 (Stochastic entropy solution). An $L^2(\mathbb{R}^d)$ -valued $\{\mathcal{F}_t : t \geq 0\}$ -predictable stochastic process $u(t) = u(t, x)$ is called a stochastic entropy solution of (1.1) if the following hold:

i) For each $T > 0$

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\|u(\cdot, t)\|_2^2 \right] < +\infty.$$

ii) For each $0 \leq \psi \in C_c^\infty([0, \infty) \times \mathbb{R}^d)$ and $\beta \in \mathcal{A}$, there holds

$$\begin{aligned} & \int_{\mathbb{R}^d} \psi(x, 0) \beta(u_0(x)) dx + \int_{\Pi_T} \left\{ \partial_t \psi(t, x) \beta(u(t, x)) + F^\beta(u(t, x)) \vec{v}(t, x) \cdot \nabla \psi(t, x) \right\} dx dt \\ & + \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \eta(u(t, x); z) \beta'(u(t, x) + \lambda \eta(u(t, x); z)) \psi(t, x) d\lambda \tilde{N}(dz, dt) dx \\ & + \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 (1 - \lambda) \eta^2(u(t, x); z) \beta''(u(t, x) + \lambda \eta(u(t, x); z)) \psi(t, x) d\lambda m(dz) dt dx \geq 0, \quad \mathbb{P} - \text{a.s.} \end{aligned}$$

Due to nonlocal nature of the Itô-Lévy formula and the missing noise-noise interaction, the Definition 2.3 does not alone give the L^1 -contraction principle in the sense of average when one tries to compare two entropy solutions directly, and hence fails to give uniqueness. For the details, we refer to see [12, 18]. However, in view of [2, 9], we can look for so called *generalized entropy solution* which are $L^2(\mathbb{R}^d \times (0, 1))$ -valued $\{\mathcal{F}_t : t \geq 0\}$ -predictable stochastic process.

Definition 2.4 (Generalized entropy solution). An $L^2(\mathbb{R}^d \times (0, 1))$ -valued $\{\mathcal{F}_t : t \geq 0\}$ -predictable stochastic process $u(t) = u(t, x, \alpha)$ is called a generalized stochastic entropy solution of (1.1) provided

(1) For each $T > 0$

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\|u(t, \cdot, \cdot)\|_2^2 \right] < +\infty.$$

(2) For all test functions $0 \leq \psi \in C_c^{1,2}([0, \infty) \times \mathbb{R}^d)$, and any $\beta \in \mathcal{A}$, the following inequality holds

$$\begin{aligned} & \int_{\mathbb{R}^d} \beta(u_0(x)) \psi(x, 0) dx + \int_{\Pi_T} \int_0^1 \left\{ \beta(u(t, x, \alpha)) \partial_t \psi(t, x) + F^\beta(u(t, x, \alpha)) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) \right\} dt dx \\ & + \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 \eta(u(t, x, \alpha); z) \beta'(u(t, x, \alpha) + \lambda \eta(u(t, x, \alpha); z)) \psi(t, x) d\alpha d\lambda \tilde{N}(dz, dt) dx \\ & + \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 (1 - \lambda) \eta^2(u(t, x, \alpha); z) \beta''(u(t, x, \alpha) + \lambda \eta(u(t, x, \alpha); z)) \\ & \quad \times \psi(t, x) d\lambda m(dz) dt dx \geq 0, \quad \mathbb{P} - \text{a.s.} \end{aligned} \tag{2.1}$$

The aim of this paper is to establish convergence of approximate solutions, constructed via flux-splitting finite volume scheme (cf. Section 3), to the unique entropy solution of (1.1), and we will do so under the following assumptions:

- A.1 $f : \mathbb{R} \mapsto \mathbb{R}$ is C^2 and Lipschitz continuous with $f(0) = 0$.
- A.2 $\vec{v} : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$ is a C^1 function with $\text{div}_x \vec{v}(t, x) = 0$ for all $(t, x) \in \Pi_T$. Furthermore, there exists $V < +\infty$ such that $|\vec{v}(t, x)| \leq V$ for all $(t, x) \in \Pi_T$.
- A.3 There exist positive constants $0 < \lambda^* < 1$ and $C > 0$, and $h_1(z) \in L^2(\mathbf{E}, m)$ with $0 \leq h_1(z) \leq 1$ such that for all $u, v \in \mathbb{R}; z \in \mathbf{E}$

$$|\eta(u; z) - \eta(v; z)| \leq \lambda^* |u - v| h_1(z); \quad |\eta(u, z)| \leq C h_1(z).$$

Moreover, $\eta(0; z) = 0$ for all $z \in \mathbf{E}$.

A.4 The initial function $u_0(x)$ is a $L^2(\mathbb{R}^d)$ -valued $\mathcal{F}_0$ measurable random variable satisfying

$$\mathbb{E}[\|u_0\|_2^2] < +\infty.$$

We have the following existence and uniqueness theorems whose proof is postponed to the Appendix.

Theorem 2.1. *Let the assumptions **A.1-A.4** hold. Then there exists a generalized entropy solution of (1.1) in the sense of Definition 2.4.*

Theorem 2.2. *Under the assumptions **A.1-A.4**, the generalized entropy solution of (1.1) is unique. Moreover, it is the unique stochastic entropy solution.*

Throughout this paper, we use the letter C to denote various generic constant which may change line to line. We denote by c_f the Lipschitz constant of f and c_η , the finite constant (which exists thanks to **A.3**) as $c_\eta = \int_{\mathbf{E}} h_1^2(z) m(dz)$. The Euclidean norm on $\mathbb{R}^d$ is denoted by $|\cdot|$.

3. FLUX-SPLITTING FINITE VOLUME SCHEME

Our main point of interest is numerical approximation for the problem (1.1). Let us first introduce the space discretization by finite volumes (control volumes). For that we need to recall the definition of so called admissible meshes for finite volume scheme (cf. [17]).

Definition 3.1 (Admissible mesh). An admissible mesh $\mathcal{T}$ of $\mathbb{R}^d$ is a family of disjoint polygonal connected subset of $\mathbb{R}^d$ satisfying the following:

- i) $\mathbb{R}^d$ is the union of the closure of the elements (called control volume) of $\mathcal{T}$.
- ii) The common interface of any two elements of $\mathcal{T}$ is included in a hyperplane of $\mathbb{R}^d$.
- iii) There exists nonnegative constant α such that

$$\begin{cases} \alpha h^d \leq |K|, \\ |\partial K| \leq \frac{1}{\alpha} h^{d-1}, \quad \forall K \in \mathcal{T}, \end{cases} \quad (3.1)$$

where $h = \sup \{ \text{diam}(K) : K \in \mathcal{T} \} < +\infty$, $|K|$ denotes the d -dimensional Lebesgue measure of K , and $|\partial K|$ represents the $(d-1)$ -dimensional Lebesgue measure of ∂K .

In the sequel, we denote the followings:

- $\mathcal{E}_K$: the set of interfaces of the control volume K .
- $\mathcal{N}(K)$: the set of control volumes neighbors of the control volume K .
- $K|L$: the common interface between K and L for any $L \in \mathcal{N}(K)$.
- $\mathcal{E}$: the set of all the interfaces of the mesh.
- $n_{K,\sigma}$: the unit normal to the interface σ , outward to the control volume K , for any $\sigma \in \mathcal{E}_K$.

Consider an admissible mesh $\mathcal{T}$ in the sense of Definition 3.1. In order to discretize the time variable, we split the time interval $[0, T]$ as follows: Let N be a positive integer and we set $\Delta t = \frac{T}{N}$. Define $t_n = n\Delta t$, $n = 0, 1, \dots, N$. Then $\{t_n : n = 0, 1, \dots, N\}$ splits the time interval $[0, T]$ into equal step length Δt .

It is well known that, the main idea behind flux-splitting finite volume method is to express a flux function f as the sum of a nondecreasing function f_1 and a non increasing function f_2 . Since the flux function f is Lipschitz continuous such a decomposition is always possible.

We propose the following flux-splitting finite volume scheme to approximate the solution of (1.1): for any $K \in \mathcal{T}$, and $n \in \{0, 1, 2, \dots, N-1\}$, we define the discrete unknowns u_K^n as follows

$$\begin{aligned} u_K^0 &= \frac{1}{|K|} \int_K u_0(x) dx, \\ \frac{|K|}{\Delta t} (u_K^{n+1} - u_K^n) &+ \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| \left\{ (v \cdot n_{K,\sigma})^+ (f_1(u_K^n) + f_2(u_L^n)) - (v \cdot n_{K,\sigma})^- (f_1(u_L^n) + f_2(u_K^n)) \right\} \\ &= \frac{|K|}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt), \end{aligned} \quad (3.2)$$

where, by denoting $d\nu$ the $d - 1$ dimensional Lebesgue measure

$$\begin{aligned}(v \cdot n_{K,\sigma})^+ &= \frac{1}{\Delta t |\sigma|} \int_{t_n}^{t_{n+1}} \int_{\sigma} (\vec{v}(t, x) \cdot n_{K,\sigma})^+ d\nu(x) dt, \\ (v \cdot n_{K,\sigma})^- &= \frac{1}{\Delta t |\sigma|} \int_{t_n}^{t_{n+1}} \int_{\sigma} (\vec{v}(t, x) \cdot n_{K,\sigma})^- d\nu(x) dt.\end{aligned}$$

Since $\operatorname{div}_x \vec{v}(t, x) = 0$ for any $(t, x) \in \Pi_T$, an elementary estimate yields

$$\sum_{\sigma \in \mathcal{E}_K} |\sigma| v \cdot n_{K,\sigma} = 0. \quad (3.3)$$

We define approximate finite volume solution on Π_T as a piecewise constant given by

$$u_{\mathcal{T},\Delta t}^h(t, x) = u_K^n \text{ for } x \in K \text{ and } t \in [t_n, t_{n+1}). \quad (3.4)$$

Remark 3.1. In view of the properties of stochastic integral with respect to compensated Poisson random measure, each u_K^n is $\mathcal{F}_{n\Delta t}$ -measurable for $K \in \mathcal{T}$ and $n \in \{0, 1, \dots, N\}$. Thus, $u_{\mathcal{T},\Delta t}^h(t, \cdot)$ is an $L^2(\mathbb{R}^d)$ -valued $\mathcal{F}_t$ -predictable stochastic process as u_0 satisfies **A.4**.

Finally, we state the main theorem of this paper.

Main Theorem. Let the assumptions **A.1-A.4** be true and $\mathcal{T}$ be an admissible mesh on $\mathbb{R}^d$ with size h in the sense of Definition 3.1. Let Δt be the time step as discuss above and assume that

$$\frac{\Delta t}{h} \rightarrow 0 \text{ as } h \rightarrow 0.$$

Let $u_{\mathcal{T},\Delta t}^h(t, x)$ be the finite volume approximation as prescribed by (3.4). Then, there exists a $L^2(\mathbb{R}^d \times (0, 1))$ -valued $\{\mathcal{F}_t : t \geq 0\}$ -predictable process $u = u(t, x, \alpha)$ such that

- i) $u(t, x, \alpha)$ is a generalized entropy solution of (1.1) and $u_{\mathcal{T},\Delta t}^h(t, x) \mapsto u(t, x, \alpha)$ in the sense of Young measure.
- (ii) $u_{\mathcal{T},\Delta t}^h(t, x) \mapsto \bar{u}(t, x)$ in $L_{\text{loc}}^p(\mathbb{R}^d; L^p(\Omega \times (0, T)))$ for $1 \leq p < 2$, where $\bar{u}(t, x) = \int_0^1 u(t, x, \alpha) d\alpha$ is the unique stochastic entropy solution of (1.1).

Remark 3.2. Under the CFL condition

$$\Delta t \leq \frac{(1 - \xi)\alpha^2 h}{c_f V}, \text{ for some } \xi \in (0, 1), \quad (3.5)$$

we have uniform moment estimate and weak BV estimate on $u_{\mathcal{T},\Delta t}^h$ for $\xi = 0$ and $\xi \in (0, 1)$ respectively (see Lemmas 4.1 and 4.2). In the deterministic case, condition (3.5) is sufficient to establish the convergence of approximate solutions to the unique entropy solution of the problem. But in the stochastic case, only this condition is not enough and hence, we assume the stronger condition, namely $\frac{\Delta t}{h} \rightarrow 0$ as $h \rightarrow 0$.

Remark 3.3. Since every Lipschitz continuous function can be expressed as the sum of nondecreasing function and a non increasing one, it suffices to prove the main theorem (cf. Theorem 3) for nondecreasing Lipschitz continuous flux function f .

For nondecreasing Lipschitz function f , the finite volume scheme (3.2) reduces to an upwind finite volume scheme

$$\begin{cases} \frac{|K|}{\Delta t} (u_K^{n+1} - u_K^n) + \sum_{\sigma \in \mathcal{E}_K} |\sigma| v \cdot n_{K,\sigma} f(u_\sigma^n) = \frac{|K|}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt), \\ u_K^0 = \frac{1}{|K|} \int_K u_0(x) dx, \end{cases} \quad (3.6)$$

where u_σ^n represents the upstream value at time t_n with respect to σ . More precisely, if σ is the interface between two control volumes K and L , then

$$u_\sigma^n = \begin{cases} u_K^n & \text{if } v \cdot n_{K,\sigma} \geq 0, \\ u_L^n & \text{if } v \cdot n_{K,\sigma} < 0. \end{cases}$$

The upwind finite volume approximate solution $u_{\mathcal{T},\Delta t}^h(t, x)$ on Π_T is defined as

$$u_{\mathcal{T},\Delta t}^h(t, x) = u_K^n \text{ for } x \in K \text{ and } t \in [t_n, t_{n+1}), \quad (3.7)$$

where the discrete unknown $u_K^n, K \in \mathcal{T}, n \in \{0, 1, \dots, N-1\}$ is computed from (3.6).

4. A PRIORI ESTIMATES

This section is devoted to *a-priori* estimates for the upwind finite volume approximate solution $u_{\mathcal{T},\Delta t}^h(t, x)$ which will be very useful to prove its convergence. We start with the following lemma which is essentially a uniform moment estimate.

Lemma 4.1. *Let $T > 0$ and the assumptions **A.1-A.4** hold. Consider an admissible mesh $\mathcal{T}$ on $\mathbb{R}^d$ with size h in the sense of Definition 3.1. Let $\Delta t = \frac{T}{N}$ be the time step for some $N \in \mathbb{N}$, satisfying the Courant-Friedrichs-Levy (CFL) condition*

$$\Delta t \leq \frac{\alpha^2 h}{c_f V}.$$

Then the upwind finite volume approximate solution $u_{\mathcal{T},\Delta t}^h$ satisfies the following bound

$$\|u_{\mathcal{T},\Delta t}^h\|_{L^\infty(0,T;L^2(\Omega \times \mathbb{R}^d))}^2 \leq e^{c_\eta T} \mathbb{E}[\|u_0\|_2^2]. \quad (4.1)$$

As a consequence, we see that $u_{\mathcal{T},\Delta t}^h$ satisfies the following bound

$$\|u_{\mathcal{T},\Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \leq T e^{c_\eta T} \mathbb{E}[\|u_0\|_2^2].$$

Proof. To prove (4.1), it is enough to prove: for $n \in \{0, 1, \dots, N-1\}$, the following property holds

$$\sum_{K \in \mathcal{T}} |K| \mathbb{E}[(u_K^n)^2] \leq (1 + \Delta t c_\eta)^n \mathbb{E}[\|u_0\|_2^2]. \quad (4.2)$$

Observe that

$$\begin{aligned} \sum_{K \in \mathcal{T}} |K| \mathbb{E}[(u_K^0)^2] &= \sum_{K \in \mathcal{T}} |K| \mathbb{E}\left[\left(\frac{1}{|K|} \int_K u_0(x) dx\right)^2\right] \\ &\leq \mathbb{E}[\|u_0\|_2^2] = (1 + \Delta t c_\eta)^0 \mathbb{E}[\|u_0\|_2^2]. \end{aligned}$$

Suppose that (4.2) holds for n . We will show that (4.2) holds for $n+1$. In view of (3.3), one has $\sum_{\sigma \in \mathcal{E}_K} |\sigma| v \cdot n_{K,\sigma} f(u_K^n) = 0$ and hence the scheme (3.6) reduces to

$$\frac{|K|}{\Delta t} (u_K^{n+1} - u_K^n) + \sum_{\sigma \in \mathcal{E}_K} |\sigma| v \cdot n_{K,\sigma} (f(u_\sigma^n) - f(u_K^n)) = \frac{|K|}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt).$$

Again, in view of the definition of u_σ^n , the above finite volume scheme is equivalent to

$$\frac{|K|}{\Delta t} (u_K^{n+1} - u_K^n) + \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_K^n) - f(u_L^n)) = \frac{|K|}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt). \quad (4.3)$$

Multiplying (4.3) by u_K^n and using the fact that $ab = \frac{1}{2}[(a+b)^2 - a^2 - b^2]$ for any $a, b \in \mathbb{R}$, we obtain

$$\begin{aligned} \frac{|K|}{2} [(u_K^{n+1})^2 - (u_K^n)^2] &= \frac{|K|}{2} (u_K^{n+1} - u_K^n)^2 - \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_K^n) - f(u_L^n)) u_K^n \\ &\quad + |K| \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) u_K^n \tilde{N}(dz, dt). \end{aligned}$$

Taking expectation, and using the fact that for any two constants $T_1, T_2 \geq 0$ with $T_1 < T_2$,

$$\mathbb{E}\left[X_{T_1} \int_{T_1}^{T_2} \int_{\mathbf{E}} \zeta(t, z) \tilde{N}(dz, dt)\right] = 0,$$

where ζ is a predictable integrand with $\mathbb{E}\left[\int_0^T \int_{\mathbf{E}} \zeta^2(t, z) m(dz) dt\right] < +\infty$ and X is an adapted process, we obtain

$$\begin{aligned} \frac{|K|}{2} \mathbb{E}\left[(u_K^{n+1})^2 - (u_K^n)^2\right] &= \frac{|K|}{2} \mathbb{E}\left[\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta^2(u_K^n; z) m(dz) dt\right] \\ &\quad + \frac{(\Delta t)^2}{2|K|} \mathbb{E}\left[\left(\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma|(v \cdot n_{K,\sigma})^-(f(u_K^n) - f(u_L^n))\right)^2\right] \\ &\quad - \Delta t \mathbb{E}\left[\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma|(v \cdot n_{K,\sigma})^-(f(u_K^n) - f(u_L^n)) u_K^n\right], \end{aligned}$$

where we have used (4.3) to replace $u_K^{n+1} - u_K^n$. Note that, thanks to (3.1), the following inequality holds

$$\frac{|\partial K|}{|K|} \leq \frac{1}{\alpha^2 h}.$$

Therefore, $\sum_{\sigma \in \mathcal{E}_K} |\sigma| |v \cdot n_{K,\sigma}| \leq V |\partial K| \leq \frac{V}{\alpha^2 h} |K|$, and hence

$$\sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- \leq \frac{V}{\alpha^2 h} |K|. \quad (4.4)$$

We use Cauchy-Schwartz inequality, the assumption **A.3** on η , and (4.4) to have

$$\begin{aligned} \frac{|K|}{2} \mathbb{E}\left[(u_K^{n+1})^2 - (u_K^n)^2\right] &\leq \Delta t \underbrace{\mathbb{E}\left[\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- \left\{ \frac{\Delta t V}{2\alpha^2 h} (f(u_K^n) - f(u_L^n))^2 - (f(u_K^n) - f(u_L^n)) u_K^n \right\}\right]}_{\mathbf{A}} \\ &\quad + \Delta t c_\eta \frac{|K|}{2} \mathbb{E}\left[(u_K^n)^2\right]. \end{aligned} \quad (4.5)$$

To estimate **A**, we use [17, Lemma 4.5] and have: for any $a, b \in \mathbb{R}$

$$b(f(b) - f(a)) \geq \phi(b) - \phi(a) + \frac{1}{2c_f} (f(b) - f(a))^2,$$

where $\phi(a) = \int_0^a s f'(s) ds$. Note that $0 \leq \phi(a) \leq c_f a^2$. Thus using the CFL condition $\frac{\Delta t V}{2\alpha^2 h} \leq \frac{1}{2c_f}$, we get

$$\mathbf{A} \leq \mathbb{E}\left[\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- (\phi(u_L^n) - \phi(u_K^n))\right]. \quad (4.6)$$

Combining (4.5) and (4.6), we obtain

$$\begin{aligned} &\frac{1}{2} \sum_{K \in \mathcal{T}} |K| \mathbb{E}\left[(u_K^{n+1})^2 - (u_K^n)^2\right] \\ &\leq \frac{\Delta t c_\eta}{2} \sum_{K \in \mathcal{T}} |K| \mathbb{E}\left[(u_K^n)^2\right] + \underbrace{\sum_{K \in \mathcal{T}} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- \mathbb{E}[\phi(u_L^n) - \phi(u_K^n)]}_{\mathbf{B}}. \end{aligned}$$

Since $\operatorname{div}_x \vec{v}(t, x) = 0$ for any $(t, x) \in \Pi_T$, one can show that $\mathbf{B} = 0$, yielding

$$\sum_{K \in \mathcal{T}} |K| \mathbb{E}\left[(u_K^{n+1})^2\right] \leq (1 + \Delta t c_\eta) \sum_{K \in \mathcal{T}} |K| \mathbb{E}\left[(u_K^n)^2\right] \leq (1 + \Delta t c_\eta)^{n+1} \mathbb{E}[|u_0|_2^2].$$

Thus (4.2) holds by mathematical induction. In other words, (4.1) holds as well. As a consequence, we have

$$\|u_{\mathcal{T},\Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t |K| \mathbb{E}[(u_K^n)^2] \leq T e^{c_\eta T} \mathbb{E}[\|u_0\|_2^2].$$

This completes the proof. $\square$

Lemma 4.2 (Weak BV estimate). *Suppose $T > 0$, and the assumptions **A.1-A.4** be true. Let $\mathcal{T}$ be an admissible mesh with size h in the sense of Definition 3.1. Let $\Delta t = \frac{T}{N}$ be the time step for some $N \in \mathbb{N}$, satisfying CFL condition*

$$\Delta t \leq \frac{(1-\xi)\alpha^2 h}{c_f V}, \text{ for some } \xi \in (0, 1). \quad (4.7)$$

Let $u_K^n : K \in \mathcal{T}, n \in \{0, 1, \dots, N-1\}$ be discrete unknowns as in (3.6). Then the followings hold:

a) There exists positive constant C , only depending on T, u_0, ξ, c_f, c_η such that

$$\sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| |v \cdot n_{K,\sigma}| \mathbb{E}[(f(u_\sigma^n) - f(u_K^n))^2] \leq C. \quad (4.8)$$

b) Let $R > 0$ be such that $h < R$. Define $\mathcal{T}_R = \{K \in \mathcal{T} : K \subset B(0, R)\}$ and $\mathcal{E}^R$ be the set of all interfaces of the mesh $\mathcal{T}_R$. Then there exists positive constant C_1 , only depending on $R, d, T, u_0, \xi, c_f, c_\eta$ such that

$$\sum_{n=0}^{N-1} \Delta t \sum_{\sigma \in \mathcal{E}^R} |\sigma| |v \cdot n_{K,\sigma}| \mathbb{E}[|f(u_\sigma^n) - f(u_K^n)|] \leq C_1 h^{-\frac{1}{2}}.$$

Proof. Multiplying (4.3) by $\Delta t u_K^n$, taking expectation and summing over $n = 0, 1, \dots, N-1$ and $K \in \mathcal{T}$, we obtain

$$\underbrace{\sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} |K| \mathbb{E}[(u_K^{n+1} - u_K^n) u_K^n]}_{\bar{\mathbf{A}}} + \underbrace{\sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t |\sigma| (v \cdot n_{K,\sigma})^- \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} \mathbb{E}[(f(u_K^n) - f(u_L^n)) u_K^n]}_{\bar{\mathbf{B}}} = 0.$$

Let us first consider $\bar{\mathbf{A}}$. Using the formula $ab = \frac{1}{2}[(a+b)^2 - a^2 - b^2]$ and (4.3), we rewrite $\bar{\mathbf{A}}$ as

$$\begin{aligned} \bar{\mathbf{A}} &= \frac{1}{2} \sum_{K \in \mathcal{T}} |K| \mathbb{E}[(u_K^N)^2 - (u_K^0)^2] - \frac{1}{2} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} |K| \mathbb{E} \left[\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta^2(u_K^n; z) m(dz) dt \right] \\ &\quad - \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \frac{(\Delta t)^2}{2|K|} \mathbb{E} \left[\left(\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_K^n) - f(u_L^n)) \right)^2 \right] \\ &\equiv \bar{\mathbf{A}}_1 + \bar{\mathbf{A}}_2 + \bar{\mathbf{A}}_3. \end{aligned}$$

Thanks to Cauchy-Schwartz inequality, CFL condition (4.7), (4.4), and assumption **A.3**

$$\begin{aligned} \bar{\mathbf{A}}_3 &\geq -\frac{1}{2} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \frac{(1-\xi)}{c_f} \mathbb{E} \left[\sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_K^n) - f(u_L^n))^2 \right], \\ \bar{\mathbf{A}}_3 &\geq -\frac{1}{2} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t |K| c_\eta \mathbb{E}[(u_K^n)^2]. \end{aligned}$$

Therefore, by using Lemma 4.1, we arrive at

$$\bar{\mathbf{A}} \geq -\frac{1}{2} T c_\eta e^{T c_\eta} \mathbb{E}[\|u_0\|_2^2] - \frac{1}{2} \mathbb{E}[\|u_0\|_2^2] - \frac{(1-\xi)}{2 c_f} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma = K|L}} |\sigma| (v \cdot n_{K,\sigma})^- \mathbb{E}[(f(u_K^n) - f(u_L^n))^2]$$

$$\geq -\frac{(1-\xi)}{2c_f} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma=K|L}} |\sigma|(v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_K^n) - f(u_L^n))^2 \right] - \frac{1}{2} \tilde{C},$$

for some constant $\tilde{C} > 0$, depending only on T, c_η, u_0 . A similar argumentations (cf. estimation of $\mathbf{A}$) reveal that

$$\bar{\mathbf{B}} \geq \frac{1}{2c_f} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma=K|L}} |\sigma|(v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_K^n) - f(u_L^n))^2 \right].$$

Since $\bar{\mathbf{A}} + \bar{\mathbf{B}} = 0$, we arrive at

$$\frac{\xi}{2c_f} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma=K|L}} |\sigma|(v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_K^n) - f(u_L^n))^2 \right] \leq \frac{1}{2} \tilde{C}$$

In other words, there exists positive constant C , only depending on T, u_0, ξ, c_f, c_η such that

$$\sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma=K|L}} |\sigma|(v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_K^n) - f(u_L^n))^2 \right] \leq C, \quad (4.9)$$

or equivalently (4.8) holds.

Let $\mathcal{T}_R = \{K \in \mathcal{T} : K \subset B(0, R)\}$ as in defined in Lemma 4.2. One can use Cauchy-Schwartz inequality along with (4.9) to have (cf. [6])

$$\sum_{K \in \mathcal{T}_R} \sum_{n=0}^{N-1} \Delta t \sum_{\substack{\sigma \in \mathcal{E}_K \\ \sigma=K|L}} |\sigma|(v \cdot n_{K,\sigma})^- \mathbb{E} \left[|f(u_K^n) - f(u_L^n)| \right] \leq C_1 h^{-\frac{1}{2}}, \quad (4.10)$$

where C_1 is a positive constant, depending only on $R, d, T, u_0, \xi, c_f, c_\eta$. Let $\mathcal{E}^R$ denotes the set of all interfaces of $\mathcal{T}_R$. Then (4.10) is equivalent to the following bound

$$\sum_{n=0}^{N-1} \Delta t \sum_{\sigma \in \mathcal{E}^R} |\sigma| |v \cdot n_{K,\sigma}| \mathbb{E} \left[|f(u_\sigma^n) - f(u_K^n)| \right] \leq C_1 h^{-\frac{1}{2}},$$

which completes the proof of the lemma. $\square$

5. ON ENTROPY INEQUALITY FOR APPROXIMATE SOLUTION

In this section, we establish entropy inequality for finite volume approximate solution. Like its deterministic counterpart, we need to consider a time-continuous approximate solution constructed from $u_{\mathcal{T}, \Delta t}^h$.

5.1. Time-continuous approximate solution. Since $\operatorname{div}_x \vec{v}(t, x) = 0$ for any $(t, x) \in \Pi_T$, the upwind finite volume scheme (3.6) can be rewritten as: for any $K \in \mathcal{T}$, and $n \in \{0, 1, \dots, N-1\}$

$$\begin{cases} u_K^{n+1} = u_K^n + \frac{\Delta t}{|K|} \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) + \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt), \\ u_K^0 = \frac{1}{|K|} \int_K u_0(x) dx. \end{cases}$$

We define a time-continuous discrete approximation, denoted by $v_K^n(\omega, s)$ on $\Omega \times [t_n, t_{n+1}]$, $n \in \{0, 1, \dots, N-1\}$ and $K \in \mathcal{T}$ from the discrete unknowns u_K^n as

$$v_K^n(\omega, s) = u_K^n + \int_{t_n}^s \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- \frac{f(u_\sigma^n) - f(u_K^n)}{|K|} + \int_{t_n}^s \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dt). \quad (5.1)$$

Note that,

$$\begin{cases} v_K^n(\omega, t_n) = u_K^n \\ v_K^n(\omega, t_{n+1}) = u_K^{n+1}. \end{cases}$$

We drop ω and write $v_K^n(\cdot)$ instead of $v_K^n(\omega, \cdot)$. Define a time-continuous approximate solution $v_{\mathcal{T}, \Delta t}^h(s, x)$ on $[0, T] \times \mathbb{R}^d$ by

$$v_{\mathcal{T}, \Delta t}^h(t, x) = v_K^n(t), \quad x \in K, \quad t \in [0, T]. \quad (5.2)$$

Next, we estimate the L^2 -error between $u_{\mathcal{T}, \Delta t}^h$ and $v_{\mathcal{T}, \Delta t}^h$. We have the following proposition.

Proposition 5.1. *Let the assumptions of Lemma 4.2 hold and $u_{\mathcal{T}, \Delta t}^h$ be the finite volume approximate solution defined by (3.6) and (3.7), and $v_{\mathcal{T}, \Delta t}^h$ be the corresponding time-continuous approximate solution prescribed by (5.1)-(5.2). Then there exist two constants $C, C_1 \in \mathbb{R}_+^*$, independent of h and Δt such that*

$$\|v_{\mathcal{T}, \Delta t}^h - u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \leq Ch + C_1 \Delta t.$$

Proof. In view of Lemmas 4.1 - 4.2, and the estimate (4.4) along with (4.7), we have

$$\begin{aligned} & \|v_{\mathcal{T}, \Delta t}^h - u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \\ &= \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \int_K \mathbb{E} \left[\int_{t_n}^s \int_{\mathbf{E}} \eta^2(u_K^n; z) m(dz) dt \right] dx ds \\ & \quad + \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \int_K \mathbb{E} \left[\left(\frac{s - \Delta t}{|K|} \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K, \sigma})^- (f(u_\sigma^n) - f(u_K^n)) \right)^2 \right] dx ds \\ & \leq \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \frac{(\Delta t)^3}{|K|} \frac{V|K|}{\alpha^2 h} \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K, \sigma})^- \mathbb{E} \left[(f(u_\sigma^n) - f(u_K^n))^2 \right] + c_\eta \Delta t \|u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \\ & \leq \frac{(\Delta t)^2 V}{\alpha^2 h} \sum_{K \in \mathcal{T}} \sum_{n=0}^{N-1} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K, \sigma})^- \mathbb{E} \left[(f(u_\sigma^n) - f(u_K^n))^2 \right] + c_\eta \Delta t \|u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \\ & \leq h \frac{(1 - \xi)^2 \alpha^2}{c_f^2 V} C + c_\eta \Delta t \|u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \leq Ch + C_1 \Delta t, \end{aligned}$$

where $C, C_1 \in \mathbb{R}_+^*$ are two constants, independent of h and Δt . This finishes the proof. $\square$

5.2. Entropy inequalities for the approximate solution. This subsection is devoted to derive the entropy inequalities for the finite volume approximate solution which will be used to prove the convergence of the numerical scheme and hence the existence of entropy solution of the underlying problem (1.1). To do so, we start with following proposition related to the entropy inequalities for the discrete unknowns u_K^n , $K \in \mathcal{T}$, $n \in \{0, 1, 2, \dots, N-1\}$.

Proposition 5.2 (Discrete entropy inequalities). *Let the assumptions **A.1-A.4** hold, and $T > 0$ be fixed. Consider an admissible mesh $\mathcal{T}$ on $\mathbb{R}^d$ with size h in the sense of Definition 3.1. Let $\Delta t = \frac{T}{N}$ be the time step for some $N \in \mathbb{N}$, satisfying*

$$\frac{\Delta t}{h} \rightarrow 0 \text{ as } h \rightarrow 0.$$

Then, $\mathbb{P}$ -a.s. in Ω , for any $\beta \in \mathcal{A}$ and for any nonnegative test function $\psi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ such that the following inequality holds:

$$\begin{aligned} & - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K (\beta(u_K^{n+1}) - \beta(u_K^n)) \psi(t_n, x) dx \\ & + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K F^\beta(u_K^n) \vec{v}(t, x) \cdot \nabla_x \psi(t_n, x) dx dt \\ & + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \beta'(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx \tilde{N}(dz, dt) \\ & + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx m(dz) dt \geq R^{h, \Delta t}, \end{aligned}$$

where $R^{h,\Delta t}$ satisfies the following condition: for any $\mathbb{P}$ -measurable set B , $\mathbb{E}[\mathbf{1}_B R^{h,\Delta t}] \rightarrow 0$ as $h \rightarrow 0$.

Proof. Let $T > 0$ be fixed and $\mathcal{T}$ be an admissible mesh on $\mathbb{R}^d$ with size h in the sense of Definition 3.1. Let $\Delta t = \frac{T}{N}$ be the time step for some $N \in \mathbb{N}$ and $t_n = n\Delta t, n \in \{0, 1, \dots, N\}$. Let $\beta \in \mathcal{A}$. Applying Itô-Lévy formula to $\beta(v_K^n)$, where v_K^n is prescribed by the equation (5.1), we have

$$\begin{aligned} \beta(v_K^n(t_{n+1})) &= \beta(v_K^n(t_n)) + \int_{t_n}^{t_{n+1}} \beta'(v_K^n(t)) \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- \frac{f(u_\sigma^n) - f(u_K^n)}{|K|} dt \\ &\quad + \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 \eta(u_K^n; z) \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) d\lambda \tilde{N}(dz, dt) \\ &\quad + \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(v_K^n(t) + \lambda \eta(u_K^n; z)) d\lambda m(dz) dt. \end{aligned} \quad (5.3)$$

Let $\psi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ be a nonnegative test function. Then there exists $R > h$ such that $\text{supp} \psi \subset [0, T] \times B(0, R - h)$. Also define $\mathcal{T}_R = \{K \in \mathcal{T} : K \subset B(0, R)\}$. We multiply the equation (5.3) by $|K|\psi_K^n$ where $\psi_K^n = \frac{1}{|K|} \int_K \psi(t_n, x) dx$ and then summing over all $K \in \mathcal{T}_R$ and $n \in \{0, 1, \dots, N-1\}$. The resulting expression reads to

$$\begin{aligned} &\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} [\beta(u_K^{n+1}) - \beta(u_K^n)] |K| \psi_K^n \\ &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \beta'(v_K^n(t)) \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \psi_K^n dt \\ &\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 \eta(u_K^n; z) \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) |K| \psi_K^n d\lambda \tilde{N}(dz, dt) \\ &\quad + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(v_K^n(t) + \lambda \eta(u_K^n; z)) |K| \psi_K^n d\lambda m(dz) dt \\ &\text{i.e. } A^{h,\Delta t} = B^{h,\Delta t} + M^{h,\Delta t} + D^{h,\Delta t}. \end{aligned} \quad (5.4)$$

Following [6], we express $B^{h,\Delta t}$ as follows.

$$B^{h,\Delta t} = B^{h,\Delta t} - B_1^{h,\Delta t} + \underbrace{B_1^{h,\Delta t} - B_2^{h,\Delta t}}_{=0} + B_2^{h,\Delta t},$$

where

$$\begin{aligned} B_1^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \beta'(u_K^n) \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \psi_K^n dt, \\ B_2^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- (F^\beta(u_\sigma^n) - F^\beta(u_K^n)) \psi_K^n dt. \end{aligned}$$

Observe that

$$B_1^{h,\Delta t} - B_2^{h,\Delta t} = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma|(v \cdot n_{K,\sigma})^- \left\{ \beta'(u_K^n) (f(u_\sigma^n) - f(u_K^n)) - (F^\beta(u_\sigma^n) - F^\beta(u_K^n)) \right\} \psi_K^n.$$

Thanks to nondecreasingness of functions f and β' , one has

$$\beta'(u_K^n) (f(u_\sigma^n) - f(u_K^n)) - (F^\beta(u_\sigma^n) - F^\beta(u_K^n)) = \int_{u_K^n}^{u_\sigma^n} (\beta'(u_K^n) - \beta'(s)) f'(s) ds \leq 0,$$

and hence $B_1^{h,\Delta t} - B_2^{h,\Delta t} \leq 0$.

By assumption **A.2**, we have $\sum_{\sigma \in \mathcal{E}_K} |\sigma| v \cdot n_{K,\sigma} F^\beta(u_K^n) \psi_K^n = 0$, and therefore

$$B_2^{h,\Delta t} = - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma}) F^\beta(u_\sigma^n) \psi_K^n.$$

Let x_σ be the center of the edge σ and ψ_σ^n be the value of $\psi(t_n, x_\sigma)$. Then,

$$\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma}) F^\beta(u_\sigma^n) \psi_\sigma^n = 0.$$

A similar argument (as described in Bauzet et al.[6, Proposition 4]) reveals that

$$B_2^{h,\Delta t} = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K F^\beta(u_K^n) \vec{v}(t, x) \cdot \nabla_x \psi(t_n, x) dx dt + R_1^{h,\Delta t} + R_2^{h,\Delta t},$$

where

$$\begin{aligned} R_1^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma}) [F^\beta(u_K^n) - F^\beta(u_\sigma^n)] (\psi_K^n - \psi_\sigma^n), \\ R_2^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} \left\{ |\sigma| (v \cdot n_{K,\sigma}) \psi_\sigma^n - \int_\sigma (v \cdot n_{K,\sigma}) \psi(t_n, x) d\nu(x) \right\} F^\beta(u_K^n). \end{aligned}$$

Combining all these, we obtain that

$$\begin{aligned} B^{h,\Delta t} &\leq \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K F^\beta(u_K^n) \vec{v}(t, x) \cdot \nabla_x \psi(t_n, x) dx dt \\ &\quad + B^{h,\Delta t} - B_1^{h,\Delta t} + R_1^{h,\Delta t} + R_2^{h,\Delta t}. \end{aligned} \tag{5.5}$$

Next we consider the term $M^{h,\Delta t}$. It can be decompose as follows:

$$M^{h,\Delta t} = M^{h,\Delta t} - M_1^{h,\Delta t} + M_1^{h,\Delta t}, \tag{5.6}$$

where

$$M_1^{h,\Delta t} = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \beta'(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx \tilde{N}(dz, dt).$$

Similarly, we rewrite $D^{h,\Delta t}$ as

$$D^{h,\Delta t} = D^{h,\Delta t} - D_1^{h,\Delta t} + D_1^{h,\Delta t}, \tag{5.7}$$

where

$$\begin{aligned} D_1^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx m(dz) dt. \end{aligned}$$

In view of (5.5), (5.6), (5.7), and (5.4) we have

$$\begin{aligned} &- \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K (\beta(u_K^{n+1}) - \beta(u_K^n)) \psi(t_n, x) dx \\ &+ \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K F^\beta(u_K^n) \vec{v}(t, x) \cdot \nabla_x \psi(t_n, x) dx dt \\ &+ \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \beta'(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx \tilde{N}(dz, dt) \end{aligned}$$

$$\begin{aligned}
& + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 (1-\lambda) \eta^2(u_K^n; z) \beta''(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx m(dz) dt \\
& \geq \underbrace{B_1^{h,\Delta t} - B^{h,\Delta t}}_{-R_1^{h,\Delta t}} - R_2^{h,\Delta t} + \underbrace{M^{h,\Delta t} - D^{h,\Delta t}}_{-R_2^{h,\Delta t}} \equiv R^{h,\Delta t}.
\end{aligned}$$

To complete the proof of the proposition, it is only required to show: for any $\mathbb{P}$ -measurable set B such that $\mathbb{E}[\mathbf{1}_B R^{h,\Delta t}] \rightarrow 0$ as $h \rightarrow 0$. Now we assume that $\frac{\Delta t}{h} \rightarrow 0$ as $h \rightarrow 0$. In this manner, with out loss of generality, we may assume that the CFL condition

$$\Delta t \leq \frac{(1-\xi)h}{c_f V} \alpha^2$$

holds for some $\xi \in (0, 1)$ and hence the estimates given in Lemmas 4.1 and 4.2 hold as well. To proceed further, we will separately show the convergence of $\mathbb{E}[\mathbf{1}_B (B_1^{h,\Delta t} - B^{h,\Delta t})]$, $\mathbb{E}[\mathbf{1}_B (M_1^{h,\Delta t} - M^{h,\Delta t})]$, $\mathbb{E}[\mathbf{1}_B (D_1^{h,\Delta t} - D^{h,\Delta t})]$, $\mathbb{E}[\mathbf{1}_B R_1^{h,\Delta t}]$, and $\mathbb{E}[\mathbf{1}_B R_2^{h,\Delta t}]$.

1. Study of $\mathbb{E}[\mathbf{1}_B (B_1^{h,\Delta t} - B^{h,\Delta t})]$: Let B be any $\mathbb{P}$ -measurable set. Then, by using (5.1) we get

$$\begin{aligned}
& \left| \mathbb{E}[\mathbf{1}_B (B_1^{h,\Delta t} - B^{h,\Delta t})] \right| \\
& = \left| \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \mathbf{1}_B \beta''(\xi_K^n) (v_K^n(s) - u_K^n) \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \psi_K^n ds \right] \right| \\
& \leq \left| \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \mathbf{1}_B \int_{t_n}^{t_{n+1}} \beta''(\xi_K^n) \frac{s - t_n}{|K|} \left(\sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \right)^2 \psi_K^n ds \right] \right| \\
& + \left| \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \mathbf{1}_B \int_{t_n}^{t_{n+1}} \beta''(\xi_K^n) \int_{t_n}^s \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dr) \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \psi_K^n ds \right] \right| \\
& \equiv T_1^{h,\Delta t} + T_2^{h,\Delta t}.
\end{aligned}$$

Following computations as in [6, estimation for $\tilde{T}_1^{h,k}$], it can be shown that

$$T_1^{h,\Delta t} \leq \frac{\Delta t}{h} \|\beta''\|_{L^\infty} \|\psi\|_{L^\infty} \frac{V}{\alpha^2} C.$$

Next, we move on to estimate $T_2^{h,\Delta t}$. Note that

$$\begin{aligned}
|T_2^{h,\Delta t}|^2 & \leq \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \left(\mathbf{1}_B \beta''(\xi_K^n) \psi_K^n \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- (f(u_\sigma^n) - f(u_K^n)) \right)^2 ds \right] \\
& \quad \times \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \left(\int_{t_n}^s \int_{\mathbf{E}} \eta(u_K^n; z) \tilde{N}(dz, dr) \right)^2 ds \right] \\
& \leq \|\beta''\|_{L^\infty}^2 \|\psi\|_{L^\infty}^2 \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \left(\sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- \right) \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_\sigma^n) - f(u_K^n))^2 \right] \\
& \quad \times \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{t_n}^s \int_{\mathbf{E}} \eta^2(u_K^n; z) m(dz) dr ds \right] \\
& \leq \frac{V}{\alpha^2 h} \|\beta''\|_{L^\infty}^2 \|\psi\|_{L^\infty}^2 \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t \sum_{\sigma \in \mathcal{E}_K} |\sigma| (v \cdot n_{K,\sigma})^- \mathbb{E} \left[(f(u_\sigma^n) - f(u_K^n))^2 \right] \\
& \quad \times c_\eta \Delta t \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} |K| \Delta t \mathbb{E}[(u_K^n)^2]
\end{aligned}$$

$$\leq \frac{\Delta t}{h} \frac{c_\eta V}{\alpha^2} \|\beta''\|_{L^\infty}^2 \|\psi\|_{L^\infty}^2 \|u_{\mathcal{T}, \Delta}\|_{L^2(\Omega \times \Pi_T)}^2 C.$$

In the above, the first inequality follows from Cauchy-Schwartz inequality, second inequality follows from Cauchy-Schwartz and BDG inequalities. In view of (4.4) and the assumption **A.3** on η , the third inequality holds true. In the last inequality, we have used the constant C given by Lemma 4.2. Here we note that the assumption $\frac{\Delta t}{h} \rightarrow 0$ as $h \rightarrow 0$ is crucial. Passing to the limit as $h \rightarrow 0$, we conclude that $\mathbb{E}[\mathbf{1}_B(B_1^{h, \Delta t} - B^{h, \Delta t})] \rightarrow 0$.

2. Study of $\mathbb{E}[\mathbf{1}_B(M_1^{h, \Delta t} - M^{h, \Delta t})]$: In view of triangle inequality, one has

$$\left| \mathbb{E}[\mathbf{1}_B(M_1^{h, \Delta t} - M^{h, \Delta t})] \right| \leq \mathcal{M}_1^{h, \Delta t} + \mathcal{M}_2^{h, \Delta t},$$

where

$$\begin{aligned} \mathcal{M}_1^{h, \Delta t} &:= \left| \mathbb{E} \left[\mathbf{1}_B \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \left\{ \beta'(u_K^n + \lambda \eta(u_K^n; z)) - \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) \right\} \right. \right. \\ &\quad \left. \left. \times (\psi(t_n, x) - \psi(t, x)) d\lambda dx \tilde{N}(dz, dt) \right] \right| \\ \mathcal{M}_2^{h, \Delta t} &:= \left| \mathbb{E} \left[\mathbf{1}_B \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \left\{ \beta'(u_K^n + \lambda \eta(u_K^n; z)) - \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) \right\} \right. \right. \\ &\quad \left. \left. \times \psi(t, x) d\lambda dx \tilde{N}(dz, dt) \right] \right|. \end{aligned}$$

Let us turn our focus on the term $\mathcal{M}_1^{h, \Delta t}$. Note that $\text{supp } \psi \subset B(0, R-h) \times [0, T]$ for some $R > h$. Using Cauchy-Schwartz inequality along with the assumptions **A.1-A.4**, and BDG inequality, we get

$$\begin{aligned} |\mathcal{M}_1^{h, \Delta t}|^2 &\leq \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \mathbb{E} \left[\left(\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 \left\{ \beta'(u_K^n + \lambda \eta(u_K^n; z)) - \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) \right\} \right. \right. \\ &\quad \left. \left. \times \eta(u_K^n; z) (\psi(t_n, x) - \psi(t, x)) d\lambda \tilde{N}(dz, dt) \right)^2 dx \right] |B(0, R)| \\ &\leq 2 \|\beta'\|_{L^\infty}^2 |B(0, R)| \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \mathbb{E} \left[\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta^2(u_K^n; z) (\psi(t_n, x) - \psi(t, x))^2 m(dz) dt \right] dx \\ &\leq 2(\Delta t)^2 \|\beta'\|_{L^\infty}^2 |B(0, R)| \|\partial_t \psi\|_{L^\infty}^2 c_\eta \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \Delta t |K| \mathbb{E}[(u_K^n)^2] \\ &\leq (\Delta t)^2 C(\beta', \partial_t \psi, c_\eta, R) \|u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \\ &\leq h^2 C(\xi, \alpha, c_f, V, \beta', \partial_t \psi, c_\eta, R) \|u_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \text{ (by (4.7)).} \end{aligned}$$

Thus, we see that

$$\mathcal{M}_1^{h, \Delta t} \longrightarrow 0 \text{ as } h \rightarrow 0.$$

Now, we estimate $\mathcal{M}_2^{h, \Delta t}$. Here we note that the boundedness of η i.e. $|\eta(u, z)| \leq Ch_1(z)$ for any $u \in \mathbb{R}$ and $z \in \mathbf{E}$ is crucial. In view of the Cauchy-Schwartz inequality and BDG inequality, we obtain

$$\begin{aligned} |\mathcal{M}_2^{h, \Delta t}|^2 &\leq \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \mathbb{E} \left[\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_0^1 \left\{ \beta'(u_K^n + \lambda \eta(u_K^n; z)) - \beta'(v_K^n(t) + \lambda \eta(u_K^n; z)) \right\}^2 \right. \\ &\quad \left. \times \eta^2(u_K^n; z) \psi^2(t, x) d\lambda m(dz) dt \right] dx |B(0, R)| \\ &\leq C(R) \|\beta''\|_{L^\infty}^2 \|\psi\|_{L^\infty}^2 \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \mathbb{E} \left[\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} |u_K^n - v_K^n(t)|^2 \eta^2(u_K^n; z) m(dz) dt \right] dx \end{aligned}$$

(by the boundedness of η)

$$\begin{aligned} &\leq C(R, \beta'', \psi,) \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \int_{t_n}^{t_{n+1}} \mathbb{E}[(u_K^n - v_K^n(t))^2] dt dx \left(\int_{\mathbf{E}} h_1^2(z) m(dz) \right) \\ &= C(R, \beta'', \psi,) c_\eta \|u_{\mathcal{T}, \Delta t}^h - v_{\mathcal{T}, \Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \longrightarrow 0 \text{ as } h \rightarrow 0, \end{aligned}$$

where in the last line, we have invoked the Proposition 5.1 and the CFL condition (4.7). Hence, $\mathbb{E}[\mathbf{1}_B(M_1^{h, \Delta t} - M^{h, \Delta t})] \rightarrow 0$ as $h \rightarrow 0$.

3. Study of $\mathbb{E}[\mathbf{1}_B(D_1^{h, \Delta t} - D^{h, \Delta t})]$: Observe that

$$\begin{aligned} &\left| \mathbb{E}[\mathbf{1}_B(D_1^{h, \Delta t} - D^{h, \Delta t})] \right| \\ &\leq \|\beta'''\|_\infty \|\psi\|_\infty \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta^2(u_K^n; z) |u_K^n - v_K^n(t)| dx m(dz) dt \right] \\ &\leq C(\beta''', \psi) \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{B(0, R)} |u_K^n - v_K^n(t)| dx dt \right] \left(\int_{\mathbf{E}} h_1^2(z) m(dz) \right) \\ &= C(\beta'', \psi, c_\eta) \|u_{\mathcal{T}, \Delta t}^h - v_{\mathcal{T}, \Delta t}^h\|_{L^1(\Omega \times B(0, R) \times [0, T])} \longrightarrow 0 \text{ as } h \rightarrow 0. \end{aligned}$$

In the above, the second inequality follows from the boundedness condition on η , and the last line holds because of Proposition 5.1 and the CFL condition (4.7).

4. Study of $\mathbb{E}[\mathbf{1}_B R_1^{h, \Delta t}]$ and $\mathbb{E}[\mathbf{1}_B R_2^{h, \Delta t}]$: Following computations as in Bauzet et al. [6, Proposition 4]) we infer that

$$\left| \mathbb{E}[\mathbf{1}_B R_1^{h, \Delta t}] \right| \leq C \|\psi_x\|_\infty \|\beta'\|_\infty \sqrt{h}; \quad \left| \mathbb{E}[\mathbf{1}_B R_2^{h, \Delta t}] \right| \leq \|\beta'\|_\infty \frac{V c_f}{\alpha^2} \bar{\varepsilon}(h) \|u_{\mathcal{T}, \Delta t}^h\|_{L^1(\Omega \times B(0, R) \times [0, T])},$$

where $\bar{\varepsilon}(r) \rightarrow 0$ as $r \rightarrow 0$.

We now combine all the above estimates to conclude: for any $\mathbb{P}$ -measurable set B such that

$$\mathbb{E}[\mathbf{1}_B R^{h, \Delta t}] \rightarrow 0 \text{ as } h \rightarrow 0.$$

This completes the proof of the proposition. $\square$

To prove convergence of the proposed scheme and hence existence of entropy solution for (1.1), one also needs a continuous entropy inequality on the discrete solutions. Regarding this, we have the following proposition which essentially gives the entropy inequality for finite volume approximate solution $u_{\mathcal{T}, \Delta t}^h$.

Proposition 5.3 (Entropy inequality for approximate solution). *Let the assumptions A.1-A.4 hold, and $T > 0$ be fixed. Let $\mathcal{T}$ be an admissible mesh on $\mathbb{R}^d$ with size h in the sense of Definition 3.1. Let $\Delta t = \frac{T}{N}$ be the time step for some $N \in \mathbb{N}$, satisfying*

$$\frac{\Delta t}{h} \rightarrow 0 \text{ as } h \rightarrow 0.$$

Then, $\mathbb{P}$ -a.s. in Ω , for any $\beta \in \mathcal{A}$ and for any nonnegative test function $\psi \in C_c^\infty([0, T] \times \mathbb{R}^d)$, the following inequality holds:

$$\begin{aligned} &\int_{\mathbb{R}^d} \beta(u_0(x)) \psi(0, x) dx + \int_{\Pi_T} \left\{ \beta(u_{\mathcal{T}, \Delta t}^h) \partial_t \psi(t, x) + F^\beta(u_{\mathcal{T}, \Delta t}^h) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) \right\} dt dx \\ &+ \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 \eta(u_{\mathcal{T}, \Delta t}^h; z) \beta'(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda \tilde{N}(dz, dt) dx \\ &+ \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 (1 - \lambda) \eta^2(u_{\mathcal{T}, \Delta t}^h; z) \beta''(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda m(dz) dt dx \\ &\geq \mathcal{R}^{h, \Delta t}, \end{aligned} \tag{5.8}$$

where for any $\mathbb{P}$ -measurable set B , $\mathbb{E}[\mathbf{1}_B \mathcal{R}^{h, \Delta t}] \rightarrow 0$ as $h \rightarrow 0$.

Proof. Let the assumptions of the proposition hold true. Since $\frac{\Delta t}{h} \rightarrow 0$ as $h \rightarrow 0$, we may assume that the CFL condition

$$\Delta t \leq \frac{(1 - \xi)h}{c_f V} \alpha^2$$

holds for some $\xi \in (0, 1)$ and hence the estimates given in Lemmas 4.1 and 4.2 and the Proposition 5.2 hold as well. Let $\psi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ be a nonnegative test function. Then there exists $R > h$ such that $\text{supp } \psi \subset [0, T] \times B(0, R - h)$. Also define $\mathcal{T}_R = \{K \in \mathcal{T} : K \subset B(0, R)\}$.

Note that $\psi(t_N, x) = 0$ for any $x \in \mathbb{R}^d$. Using the summation by parts formula,

$$\sum_{n=1}^N a_n (b_n - b_{n-1}) = a_N b_N - a_0 b_0 - \sum_{n=0}^{N-1} b_n (a_{n+1} - a_n)$$

one has

$$\begin{aligned} & - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K (\beta(u_K^{n+1}) - \beta(u_K^n)) \psi(t_n, x) dx \\ & = \sum_{K \in \mathcal{T}_R} \int_K \beta(u_K^0) \psi(0, x) dx + \int_{\Delta t}^T \int_{\mathbb{R}^d} \beta(u_{\mathcal{T}, \Delta t}^h) \partial_t \psi(t - \Delta t, x) dx dt. \end{aligned} \quad (5.9)$$

Let $R^{h, \Delta t}$ be the quantity as in the Proposition 5.2. Define

$$\begin{aligned} \mathcal{R}^{h, \Delta t} &= R^{h, \Delta t} + \left\{ \int_{\mathbb{R}^d} \beta(u_0(x)) \psi(0, x) dx - \sum_{K \in \mathcal{T}_R} \int_K \beta(u_K^0) \psi(0, x) dx \right\} \\ &+ \left\{ \int_{\Pi_T} \beta(u_{\mathcal{T}, \Delta t}) \partial_t \psi(t, x) dt dx - \int_{\Delta t}^T \int_{\mathbb{R}^d} \beta(u_{\mathcal{T}, \Delta t}^h) \partial_t \psi(t - \Delta t, x) dx dt \right\} \\ &+ \left\{ \int_{\Pi_T} F^\beta(u_{\mathcal{T}, \Delta t}^h) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) dt dx - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K F^\beta(u_K^n) \vec{v}(t, x) \cdot \nabla_x \psi(t_n, x) dx dt \right\} \\ &+ \left\{ \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 \eta(u_{\mathcal{T}, \Delta t}^h; z) \beta'(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda \tilde{N}(dz, dt) dx \right. \\ &\quad \left. - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta(u_K^n; z) \beta'(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx \tilde{N}(dz, dt) \right\} \\ &+ \left\{ \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 (1 - \lambda) \eta^2(u_{\mathcal{T}, \Delta t}^h; z) \beta''(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda m(dz) dt dx \right. \\ &\quad \left. - \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(u_K^n + \lambda \eta(u_K^n; z)) \psi(t_n, x) d\lambda dx m(dz) dt \right\} \\ &\equiv R^{h, \Delta t} + \mathcal{I}^{h, \Delta t} + \mathcal{T}^{h, \Delta t} + \mathcal{D}^{h, \Delta t} + \mathcal{M}^{h, \Delta t} + \mathcal{A}^{h, \Delta t}. \end{aligned}$$

In view of Proposition 5.2 and the definition of $\mathcal{R}^{h, \Delta t}$ along with (5.9), we note that (5.8) holds.

In order to prove the proposition, it remains to prove the convergence of the following quantities: $\mathbb{E}[\mathbf{1}_B R^{h, \Delta t}]$, $\mathbb{E}[\mathbf{1}_B \mathcal{I}^{h, \Delta t}]$, $\mathbb{E}[\mathbf{1}_B \mathcal{T}^{h, \Delta t}]$, $\mathbb{E}[\mathbf{1}_B \mathcal{D}^{h, \Delta t}]$, $\mathbb{E}[\mathbf{1}_B \mathcal{M}^{h, \Delta t}]$ and $\mathbb{E}[\mathbf{1}_B \mathcal{A}^{h, \Delta t}]$, where B is any $\mathbb{P}$ -measurable subset of Ω .

1. **Convergence of $\mathbb{E}[\mathbf{1}_B \mathcal{I}^{h,\Delta t}]$** : Note that, due to Lebesgue differentiation theorem, for almost all $x \in K$, $|u_0(x) - u_K^0| \rightarrow 0$ as diameter of K tends to zero (i.e., $h \rightarrow 0$). Now

$$\begin{aligned} \left| \mathbb{E}[\mathbf{1}_B \mathcal{I}^{h,\Delta t}] \right| &= \left| \mathbb{E} \left[\mathbf{1}_B \sum_{K \in \mathcal{T}_R} \int_K (\beta(u_0(x)) - \beta(u_K^0)) \psi(x, 0) dx \right] \right| \\ &\leq \|\beta'\|_\infty \mathbb{E} \left[\sum_{K \in \mathcal{T}_R} \int_K |u_0(x) - u_K^0| \psi(x, 0) dx \right], \end{aligned}$$

and hence $\mathbb{E}[\mathbf{1}_B \mathcal{I}^{h,\Delta t}] \rightarrow 0$ as $h \rightarrow 0$.

2. **Convergence of $\mathbb{E}[\mathbf{1}_B \mathcal{T}^{h,\Delta t}]$** : In view of Lemma 4.2 and CFL condition (4.7), we obtain

$$\begin{aligned} &\left| \mathbb{E}[\mathbf{1}_B \mathcal{T}^{h,\Delta t}] \right| \\ &= \left| \mathbb{E} \left[\mathbf{1}_B \int_0^{\Delta t} \int_{\mathbb{R}^d} \beta(u_{\mathcal{T},\Delta t}^h) \partial_t \psi(x, t) dx dt \right] + \mathbb{E} \left[\mathbf{1}_B \int_{\Delta t}^T \int_{\mathbb{R}^d} \beta(u_{\mathcal{T},\Delta t}^h) (\partial_t \psi(t, x) - \partial_t \psi(t - \Delta t, x)) dx dt \right] \right| \\ &\leq \|\beta'\|_\infty \Delta t \left(\|\partial_t \psi\|_\infty \|u_{\mathcal{T},\Delta t}^h\|_{L^\infty(0,T;L^1(\Omega \times B(0,R)))} + \|\partial_{tt} \psi\|_\infty \|u_{\mathcal{T},\Delta t}^h\|_{L^1(\Omega \times B(0,R) \times [0,T])} \right), \end{aligned}$$

and hence $\mathbb{E}[\mathbf{1}_B \mathcal{T}^{h,\Delta t}] \rightarrow 0$ as $h \rightarrow 0$.

3. **Convergence of $\mathbb{E}[\mathbf{1}_B \mathcal{D}^{h,\Delta t}]$** : In view of A.3, one has

$$\begin{aligned} \left| \mathbb{E}[\mathbf{1}_B \mathcal{D}^{h,\Delta t}] \right| &\leq V \|\nabla_x \partial_t \psi\|_\infty \Delta t \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \int_{t_n}^{t_{n+1}} \mathbb{E} \left[|F^\beta(u_K^n)| \right] dx dt \\ &\leq V \|\nabla_x \partial_t \psi\|_\infty \Delta t \|\beta'\|_\infty c_f \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_K \int_{t_n}^{t_{n+1}} \mathbb{E} [|u_K^n|] dx dt \\ &\leq V \|\nabla_x \partial_t \psi\|_\infty \Delta t \|\beta'\|_\infty c_f \|u_{\mathcal{T},\Delta t}^h\|_{L^1(\Omega \times B(0,R) \times [0,T])} \rightarrow 0 \text{ as } h \rightarrow 0. \end{aligned}$$

Thanks to the Lemma 4.1 and the CFL condition (4.7), one can pass to the limit in the last line as well.

4. **Convergence of $\mathbb{E}[\mathbf{1}_B \mathcal{M}^{h,\Delta t}]$** : By using Cauchy-Schwartz inequality, BDG inequality, CFL condition (4.7) and Lemma 4.1, we obtain

$$\begin{aligned} \left| \mathbb{E}[\mathbf{1}_B \mathcal{M}^{h,\Delta t}] \right|^2 &\leq |B(0, R)| \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 \eta^2(u_K^n; z) \beta'^2(u_K^n + \lambda \eta(u_K^n; z)) \right. \\ &\quad \left. \times (\psi(t, x) - \psi(t_n, x))^2 d\lambda dx m(dz) dt \right] \\ &\leq C(R, \psi, c_\eta) (\Delta t)^2 \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K \mathbb{E} [(u_K^n)^2] dx dt \\ &\leq C(R, \psi, c_\eta) (\Delta t)^2 \|u_{\mathcal{T},\Delta t}^h\|_{L^2(\Omega \times \Pi_T)}^2 \rightarrow 0 \text{ as } h \rightarrow 0. \end{aligned}$$

5. **Convergence of $\mathbb{E}[\mathbf{1}_B \mathcal{A}^{h,\Delta t}]$** : Note that

$$\begin{aligned} \mathcal{A}^{h,\Delta t} &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \int_0^1 (1 - \lambda) \eta^2(u_K^n; z) \beta''(u_K^n + \lambda \eta(u_K^n; z)) \\ &\quad \times \left\{ \psi(t, x) - \psi(t_n, x) \right\} d\lambda dx m(dz) dt. \end{aligned}$$

Therefore, by (4.7) and Lemma 4.1, we obtain

$$\left| \mathbb{E}[\mathbf{1}_B \mathcal{A}^{h,\Delta t}] \right| \leq \|\beta''\|_\infty \Delta t \|\partial_t \psi\|_\infty \mathbb{E} \left[\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \int_K \eta^2(u_K^n; z) dx m(dz) dt \right]$$

$$\leq C(\beta, \psi, c_\eta) \Delta t \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}_R} \int_{t_n}^{t_{n+1}} \int_K \mathbb{E}[(u_K^n)^2] dx dt \longrightarrow 0 \text{ as } h \rightarrow 0.$$

6. Convergence of $\mathbb{E}[\mathbf{1}_B R^{h, \Delta t}]$: Thanks to Proposition 5.2, we have seen that

$$\mathbb{E}[\mathbf{1}_B R^{h, \Delta t}] \longrightarrow 0 \text{ as } h \rightarrow 0,$$

for any $\mathbb{P}$ -measurable set B . □

6. PROOF OF THE MAIN THEOREM

In this section, we establish the convergence of the scheme and hence existence of entropy solution to the underlying problem (1.1). Note that *a-priori* estimates on $u_{\mathcal{T}, \Delta t}^h(x, t)$ given by Lemma 4.1 only guarantee weak compactness of the family $\{u_{\mathcal{T}, \Delta t}^h\}_{h>0}$, which is inadequate in view of the nonlinearities in the equation. The concept of Young measure theory is appropriate in this case. We now recapitulate the results we shall use from Young measure theory due to Dafermos [13] and Panov [28] for the deterministic setting, and Balder [1] for the stochastic version of the theory.

6.1. Young measure and convergence of approximate solutions. Roughly speaking a Young measure is a parametrized family of probability measures where the parameters are drawn from a measure space. Let (Θ, Σ, μ) be a σ -finite measure space and $\mathcal{P}(\mathbb{R})$ be the space of probability measures on $\mathbb{R}$.

Definition 6.1 (Young Measure). A Young measure from Θ into $\mathbb{R}$ is a map $\tau \mapsto \mathcal{P}(\mathbb{R})$ such that $\tau(\cdot) : \theta \mapsto \tau(\theta)(B)$ is Σ -measurable for every Borel subset B of $\mathbb{R}$. The set of all Young measures from Θ into $\mathbb{R}$ is denoted by $\mathcal{R}(\Theta, \Sigma, \mu)$.

In this context, we mention that with an appropriate choice of (Θ, Σ, μ) , the family $\{u_{\mathcal{T}, \Delta t}^h\}_{h>0}$ can be thought of as a family of Young measures. We are interested in finding a subsequences out of this family that “converges” to a Young measure in a suitable sense. We sum up the necessary results in the following lemma to carry over the subsequent analysis. For a proof of this lemma, consult [2, 9].

Proposition 6.1. *Let $\{u_{\mathcal{T}, \Delta t}^h(t, x)\}_{h>0}$ be a sequence of $L^2(\mathbb{R}^d)$ -valued predictable processes such that (4.1) holds. Then there exists a subsequence $\{h_n\}$ with $h_n \rightarrow 0$ and a Young measure $\tau \in \mathcal{R}(\Theta, \Sigma, \mu)$ such that the following hold:*

- (A) *If $g(\theta, \xi)$ is a Caratheodory function on $\Theta \times \mathbb{R}$ such that $\text{supp}(g) \subset \Theta_M \times \mathbb{R}$ for some $M \in \mathbb{N}$ and $\{g(\theta, u_{\mathcal{T}, \Delta t}^{h_n}(\theta))\}_n$ (where $\theta \equiv (\omega; t, x)$) is uniformly integrable, then*

$$\lim_{h_n \rightarrow 0} \int_{\Theta} g(\theta, u_{\mathcal{T}, \Delta t}^{h_n}(\theta)) \mu(d\theta) = \int_{\Theta} \left[\int_{\mathbb{R}} g(\theta, \xi) \tau(\theta)(d\xi) \right] \mu(d\theta).$$

- (B) *Denoting a triplet $(\omega, x, t) \in \Theta$ by θ , we define*

$$u(\theta, \alpha) = \inf \left\{ c \in \mathbb{R} : \tau(\theta)((-\infty, c)) > \alpha \right\} \quad \text{for } \alpha \in (0, 1) \text{ and } \theta \in \Theta.$$

Then, $u(\theta, \alpha)$ is non-decreasing, right continuous on $(0, 1)$ and $\mathcal{P}_T \times \mathcal{L}(\mathbb{R}^d \times (0, 1))$ -measurable. Moreover, if $g(\theta, \xi)$ is a nonnegative Caratheodory function on $\Theta \times \mathbb{R}$, then

$$\int_{\Theta} \left[\int_{\mathbb{R}} g(\theta, \xi) \tau(\theta)(d\xi) \right] \mu(d\theta) = \int_{\Theta} \int_{\alpha=0}^1 g(\theta, u(\theta, \alpha)) d\alpha \mu(d\theta).$$

6.2. Proof of the main theorem. Having all the necessary *a-priori* bounds and entropy inequality on $u_{\mathcal{T}, \Delta t}^h$, we are now ready to prove the main theorem (cf. Main Theorem 3). Here we mentioned that $u(\theta, \alpha)$ given by Proposition 6.1 will serve as a possible generalized entropy solution to (1.1) for a specific choice of measure space (Θ, Σ, μ) .

In view of (5.8), we have for any $B \in \mathcal{F}_T$

$$\begin{aligned} & \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \left\{ \beta(u_{\mathcal{T}, \Delta t}^h) \partial_t \psi(t, x) + F^\beta(u_{\mathcal{T}, \Delta t}^h) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) \right\} dt dx \right] \\ & + \mathbb{E} \left[\mathbf{1}_B \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 \eta(u_{\mathcal{T}, \Delta t}^h; z) \beta'(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda \tilde{N}(dz, dt) dx \right] \end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \left[\mathbf{1}_B \int_{\mathbb{R}^d} \int_0^T \int_{\mathbf{E}} \int_0^1 (1-\lambda) \eta^2(u_{\mathcal{T}, \Delta t}^h; z) \beta''(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda m(dz) dt dx \right] \\
& + \mathbb{E} \left[\mathbf{1}_B \int_{\mathbb{R}^d} \beta(u_0(x)) \psi(0, x) dx \right] \geq \mathbb{E} \left[\mathbf{1}_B \mathcal{R}^{h, \Delta t} \right].
\end{aligned} \tag{6.1}$$

We would like to pass the limit in (6.1) as h approaches to zero. To do this, here we use the technique of Young measure theory in stochastic setting. Let the predictable σ -field of $\Omega \times (0, T)$ with the respect to $\{\mathcal{F}_t\}$ is denoted by $\mathcal{P}_T$, and we set

$$\Theta = \Omega \times (0, T) \times \mathbb{R}^d, \quad \Sigma = \mathcal{P}_T \times \mathcal{L}(\mathbb{R}^d) \quad \text{and} \quad \mu = P \otimes \lambda_t \otimes \lambda_x,$$

where λ_t and λ_x are respectively the Lebesgue measures on $(0, T)$ and $\mathbb{R}^d$. Moreover, for $M \in \mathbb{N}$, set $\Theta_M = \Omega \times (0, T) \times B_M$, where B_M be the ball of radius M around zero in $\mathbb{R}^d$.

In view of the Proposition 6.1 one can show (see the Corollary 4.6 of [9]) that for every $B \in \mathcal{F}_T$

$$\begin{aligned}
& \lim_{h \rightarrow 0} \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \eta(u_{\mathcal{T}, \Delta t}^h; z) \beta'(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda \tilde{N}(dz, dt) dx \right] \\
& = \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 \eta(u(x, t, \alpha); z) \beta'(u(x, t, \alpha) + \lambda \eta(u(x, t, \alpha); z)) \psi(t, x) d\alpha d\lambda \tilde{N}(dz, dt) dx \right].
\end{aligned} \tag{6.2}$$

Note that for any $B \in \mathcal{F}_T$, the function $\mathbf{1}_B \in L^2\left(0, T; L^2((\Omega, \mathcal{F}_T), L^2(\mathbb{R}))\right)$, and $L^2(\Theta, \Sigma, \mu)$ is closed subspace of $L^2\left(0, T; L^2((\Omega, \mathcal{F}_T), L^2(\mathbb{R}))\right)$. Hence the weak convergence in $L^2(\Theta, \Sigma, \mu)$ would imply weak convergence in $L^2\left(0, T; L^2((\Omega, \mathcal{F}_T), L^2(\mathbb{R}^d))\right)$. Therefore, in view of Proposition 6.1, we have

$$\begin{aligned}
& \lim_{h \rightarrow 0} \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \left\{ \beta(u_{\mathcal{T}, \Delta t}^h) \partial_t \psi(t, x) + F^\beta(u_{\mathcal{T}, \Delta t}^h) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) \right\} dt dx \right] \\
& = \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \left\{ \beta(u(x, t, \alpha)) \partial_t \psi(t, x) + F^\beta(u(x, t, \alpha)) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) \right\} d\alpha dt dx \right]
\end{aligned} \tag{6.3}$$

and

$$\begin{aligned}
& \lim_{h \rightarrow 0} \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 (1-\lambda) \eta^2(u_{\mathcal{T}, \Delta t}^h; z) \beta''(u_{\mathcal{T}, \Delta t}^h + \lambda \eta(u_{\mathcal{T}, \Delta t}^h; z)) \psi(t, x) d\lambda m(dz) dt dx \right] \\
& = \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 (1-\lambda) \eta^2(u(x, t, \alpha); z) \beta''(u(x, t, \alpha) + \lambda \eta(u(x, t, \alpha); z)) \psi(t, x) d\alpha d\lambda m(dz) dt dx \right].
\end{aligned} \tag{6.4}$$

By (6.3), (6.4) and (6.2) and the fact that $\mathbb{E} \left[\mathbf{1}_B \mathcal{R}^{h, \Delta t} \right] \rightarrow 0$ as $h \rightarrow 0$ (cf. Proposition 5.3), one can pass to the limit in (6.1) yielding (2.1). Also, in view of Proposition 6.1 and the uniform moment estimate (4.1) along with Fatou's lemma, we have

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\|u(t, \cdot, \cdot)\|_2^2 \right] < +\infty.$$

This implies that $u(t, x, \alpha)$ is a generalized entropy solution of (1.1). Again, thanks to Theorem 2.2, we conclude that $u(t, x, \alpha)$ is an independent function of variable α and $\bar{u}(t, x) = \int_0^1 u(t, x, \tau) d\tau = u(t, x, \alpha)$ (for all α) is the unique stochastic entropy solution. Moreover, since $u_{\mathcal{T}, \Delta t}^h$ is bounded in $L^2(\Omega \times \Pi_T)$, we conclude that $u_{\mathcal{T}, \Delta t}^h$ converges to $\bar{u}$ in $L_{\text{loc}}^p(\mathbb{R}^d; L^p(\Omega \times (0, T)))$, for $1 \leq p < 2$. This completes the proof.

7. APPENDIX

In this section, we study existence and uniqueness of entropy solution for the underlying problem (1.1).

7.1. Existence of weak solution for viscous problem. Just as the deterministic problem, here also we study the corresponding regularized problem by adding a small diffusion operator and derive some *a-priori* bounds. Due to the nonlinearity in equation, one cannot expect classical solution and instead seeks an weak solution.

For a small parameter $\varepsilon > 0$, we consider the following viscous approximation of (1.1)

$$\begin{aligned} du(t, x) + \operatorname{div}_x(\vec{v}(t, x)f(u(t, x))) dt &= \int_{\mathbf{E}} \eta(u(t, x); z) \tilde{N}(dz, dt) + \varepsilon \Delta u(t, x) dt, \quad (t, x) \in \Pi_T \\ u(0, x) &= u_0^\varepsilon(x), \quad x \in \mathbb{R}^d. \end{aligned} \quad (7.1)$$

To establish existence of a weak solution for (7.1), we follow [11, 30] and use an implicit time discretization scheme. Let $\Delta t = \frac{T}{N}$ for some fixed positive integer $N \geq 1$. Set $t_n = n \Delta t$ for $n = 0, 1, 2, \dots, N$. Define

$$\begin{aligned} \mathcal{N} &= L^2(\Omega; H^1(\mathbb{R}^d)), \quad \mathcal{N}_n = \{\text{the } \mathcal{F}_{n\Delta t} \text{ measurable elements of } \mathcal{N}\}, \\ \mathcal{H} &= L^2(\Omega; L^2(\mathbb{R}^d)), \quad \mathcal{H}_n = \{\text{the } \mathcal{F}_{n\Delta t} \text{ measurable elements of } \mathcal{H}\}. \end{aligned}$$

The following proposition holds.

Proposition 7.1. *Assume that Δt is small with $\Delta t < \frac{2\varepsilon}{V^2 c_f^2}$. Then, for any given $u_n \in \mathcal{H}_n$, there exists a unique $u_{n+1} \in \mathcal{N}_{n+1}$ such that $\mathbb{P}$ -a.s. for any $v \in H^1(\mathbb{R}^d)$, the following variational formula holds:*

$$\begin{aligned} &\int_{\mathbb{R}^d} \left((u_{n+1} - u_n)v + \Delta t \{ \varepsilon \nabla u_{n+1} \cdot \nabla v - \vec{v}(t_n, x)f(u_{n+1}) \cdot \nabla v \} \right) dx \\ &= \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) v \tilde{N}(dz, ds) dx. \end{aligned} \quad (7.2)$$

Proof. Let us define a map

$$\begin{aligned} T : \mathcal{H}_{n+1} &\mapsto \mathcal{H}_{n+1} \\ S &\mapsto u = T(S) \end{aligned}$$

via the variational problem in $\mathcal{N}_{n+1}$: for all $v \in \mathcal{N}_{n+1}$

$$\begin{aligned} &\mathbb{E} \left[\int_{\mathbb{R}^d} \left((u - u_n)v + \Delta t \{ \varepsilon \nabla u \cdot \nabla v - \vec{v}(t_n, x)f(S) \cdot \nabla v \} \right) dx \right] \\ &= \mathbb{E} \left[\int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) v \tilde{N}(dz, ds) dx \right]. \end{aligned}$$

Thanks to Lax-Milgram theorem, T is a well-defined function. Moreover, for any $S_1, S_2 \in \mathcal{H}_{n+1}$, we see that

$$\begin{aligned} &\mathbb{E} \left[\int_{\mathbb{R}^d} |T(S_1) - T(S_2)|^2 dx + \Delta t \varepsilon \int_{\mathbb{R}^d} |\nabla(T(S_1) - T(S_2))|^2 dx \right] \\ &= \Delta t \mathbb{E} \left[\int_{\mathbb{R}^d} \vec{v}(t_n, x)(f(S_1) - f(S_2)) \cdot \nabla(T(S_1) - T(S_2)) dx \right] \end{aligned}$$

and hence, by Young's inequality and assumptions **A.1** and **A.2**

$$\begin{aligned} &\mathbb{E} \left[\int_{\mathbb{R}^d} |T(S_1) - T(S_2)|^2 dx + \frac{\Delta t}{2} \varepsilon \int_{\mathbb{R}^d} |\nabla(T(S_1) - T(S_2))|^2 dx \right] \\ &\leq \frac{\Delta t}{2\varepsilon} \mathbb{E} \left[\int_{\mathbb{R}^d} |\vec{v}(t_n, x)|^2 |f(S_1) - f(S_2)|^2 dx \right] \leq \frac{\Delta t V^2 c_f^2}{2\varepsilon} \mathbb{E} \left[\int_{\mathbb{R}^d} |S_1 - S_2|^2 dx \right] \end{aligned}$$

Thus, if $\Delta t < \frac{2\varepsilon}{V^2 c_f^2}$, then T is a contractive mapping in $\mathcal{H}_{n+1}$ which completes the proof. $\square$

7.1.1. A-priori estimate. Note that, since $\operatorname{div}_x \vec{v}(t, x) = 0$ for all $(t, x) \in \Pi_T$, for any $\theta \in \mathcal{D}(\mathbb{R}^d)$, $\int_{\mathbb{R}^d} \vec{v}(t, x) f(\theta) \cdot \nabla \theta dx = 0$ and hence true for any $\theta \in H^1(\mathbb{R}^d)$ by density argument. We choose a test function $v = u_{n+1}$ in (7.2) and have

$$\begin{aligned} & \int_{\mathbb{R}^d} (u_{n+1} - u_n) u_{n+1} dx + \varepsilon \Delta t \int_{\mathbb{R}^d} |\nabla u_{n+1}|^2 dx = \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds) u_{n+1} dx \\ & \leq \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) u_n \tilde{N}(dz, ds) dx + \frac{\alpha}{2} \|u_{n+1} - u_n\|_{L^2(\mathbb{R}^d)}^2 \\ & \quad + \frac{1}{2\alpha} \int_{\mathbb{R}^d} \left(\int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds) \right)^2 dx, \quad \text{for some } \alpha > 0. \end{aligned} \quad (7.3)$$

Therefore, thanks to the assumption **A.3**, and Itô-Lévy isometry

$$\frac{1}{2} \left[\|u_{n+1}\|_{\mathcal{H}}^2 + \|u_{n+1} - u_n\|_{\mathcal{H}}^2 - \|u_n\|_{\mathcal{H}}^2 \right] + \varepsilon \Delta t \|\nabla u_{n+1}\|_{\mathcal{H}}^2 \leq \frac{\alpha}{2} \|u_{n+1} - u_n\|_{\mathcal{H}}^2 + \frac{C \Delta t}{2\alpha} (1 + \|u_n\|_{\mathcal{H}}^2).$$

Since $\alpha > 0$ is arbitrary, one can choose $\alpha > 0$ so that

$$\|u_n\|_{\mathcal{H}}^2 + \sum_{k=0}^{n-1} \|u_{k+1} - u_k\|_{\mathcal{H}}^2 + \varepsilon \Delta t \sum_{k=0}^{n-1} \|\nabla u_{k+1}\|_{\mathcal{H}}^2 \leq C_1 + C_2 \Delta t \sum_{k=0}^{n-1} \|u_k\|_{\mathcal{H}}^2,$$

for some constants $C_1, C_2 > 0$. Hence an application of discrete Gronwall's lemma implies

$$\|u_n\|_{\mathcal{H}}^2 + \sum_{k=0}^{n-1} \|u_{k+1} - u_k\|_{\mathcal{H}}^2 + \varepsilon \Delta t \sum_{k=0}^{n-1} \|\nabla u_{k+1}\|_{\mathcal{H}}^2 \leq C. \quad (7.4)$$

For fixed $\Delta t = \frac{T}{N}$, we define

$$u^{\Delta t}(t) = \sum_{k=1}^N u_k \mathbf{1}_{[t_{k-1}, t_k)}(t); \quad \tilde{u}^{\Delta t}(t) = \sum_{k=1}^N \left[\frac{u_k - u_{k-1}}{\Delta t} (t - t_{k-1}) + u_{k-1} \right] \mathbf{1}_{[t_{k-1}, t_k)}(t)$$

with $u^{\Delta t}(t) = u_0$ for $t < 0$. Similarly, we define

$$\tilde{B}^{\Delta t}(t) = \sum_{k=1}^N \left[\frac{B_k - B_{k-1}}{\Delta t} (t - t_{k-1}) + B_{k-1} \right] \mathbf{1}_{[t_{k-1}, t_k)}(t),$$

where

$$B_n = \sum_{k=0}^{n-1} \int_{t_k}^{t_{k+1}} \int_{\mathbf{E}} \eta(u_k; z) \tilde{N}(dz, ds) = \int_0^{t_n} \int_{\mathbf{E}} \eta(u^{\Delta t}(s - \Delta t); z) \tilde{N}(dz, ds).$$

A straightforward calculation shows that

$$\begin{cases} \|u^{\Delta t}\|_{L^\infty(0, T; \mathcal{H})} = \max_{k=1, 2, \dots, N} \|u_k\|_{\mathcal{H}}; & \|\tilde{u}^{\Delta t}\|_{L^\infty(0, T; \mathcal{H})} = \max_{k=0, 1, \dots, N} \|u_k\|_{\mathcal{H}}, \\ \|u^{\Delta t} - \tilde{u}^{\Delta t}\|_{L^2(0, T; \mathcal{H})}^2 \leq \Delta t \sum_{k=0}^{N-1} \|u_{k+1} - u_k\|_{\mathcal{H}}^2. \end{cases}$$

In view of the above definitions and *a-priori* estimate (7.4), we have the following lemma.

Lemma 7.2. *Assume that Δt is small. Then $u^{\Delta t}, \tilde{u}^{\Delta t}$ are bounded sequences in $L^\infty(0, T; \mathcal{H})$; $\sqrt{\varepsilon} u^{\Delta t}$ is a bounded sequence in $L^2(0, T; \mathcal{N})$ and $\|u^{\Delta t} - \tilde{u}^{\Delta t}\|_{L^2(0, T; \mathcal{H})}^2 \leq C \Delta t$. Moreover, $u^{\Delta t} - u^{\Delta t}(\cdot - \Delta t) \rightarrow 0$ in $L^2(\Omega \times \Pi_T)$.*

Next, we want to find some upper bound for $\tilde{B}^{\Delta t}(t)$. Regarding this, we have the following lemma.

Lemma 7.3. *$\tilde{B}^{\Delta t}$ is a bounded sequence in $L^2(\Omega \times \Pi_T)$ and*

$$\left\| \tilde{B}^{\Delta t}(\cdot) - \int_0^\cdot \int_{\mathbf{E}} \eta(u^{\Delta t}(s - \Delta t); z) \tilde{N}(dz, ds) \right\|_{L^2(\Omega \times \mathbb{R}^d)}^2 \leq C \Delta t.$$

Proof. First we prove the boundedness of $\tilde{B}^{\Delta t}(t)$. By using definition of $\tilde{B}^{\Delta t}(t)$, the assumption **A.3**, and boundedness of $u^{\Delta t}$ in $L^\infty(0, T; \mathcal{H})$, we obtain

$$\begin{aligned} \|\tilde{B}^{\Delta t}\|_{L^2(0, T; L^2(\Omega, L^2(\mathbb{R}^d)))}^2 &\leq \Delta t \sum_{k=0}^N \|B_k\|_{L^2(\Omega \times \mathbb{R}^d)}^2 \\ &\leq \Delta t \sum_{k=0}^N \mathbb{E} \left[\left| \int_{\mathbb{R}^d} \int_0^{t_k} \int_{\mathbf{E}} \eta(u^{\Delta t}(s - \Delta t); z) \tilde{N}(dz, ds) dx \right|^2 \right] \\ &\leq C \Delta t \sum_{k=0}^N \mathbb{E} \left[\int_{\mathbb{R}^d} \int_0^{t_k} |u^{\Delta t}(s - \Delta t)|^2 dx ds \right] \leq C \|u^{\Delta t}\|_{L^\infty(0, T; L^2(\Omega \times \mathbb{R}^d))} \leq C. \end{aligned}$$

Thus, $\tilde{B}^{\Delta t}$ is a bounded sequence in $L^2(\Omega \times \Pi_T)$.

To prove second part of the lemma, we see that for any $t \in [t_n, t_{n+1})$,

$$\begin{aligned} \tilde{B}^{\Delta t}(t) &= \int_0^t \int_{\mathbf{E}} \eta(u^{\Delta t}(s - \Delta t); z) \tilde{N}(dz, ds) \\ &= \frac{t - t_n}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds) - \int_{t_n}^t \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds). \end{aligned}$$

Therefore, in view of (7.4) and assumption **A.3**, we have

$$\begin{aligned} &\left\| \tilde{B}^{\Delta t}(t) - \int_0^t \int_{\mathbf{E}} \eta(u^{\Delta t}(s - \Delta t); z) \tilde{N}(dz, ds) \right\|_{L^2(\Omega \times \mathbb{R}^d)}^2 \\ &\leq 2 \int_{\mathbb{R}^d} \mathbb{E} \left[\left(\frac{t - t_n}{\Delta t} \right)^2 \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta^2(u_n; z) m(dz) ds + \int_{t_n}^t \int_{\mathbf{E}} \eta^2(u_n; z) m(dz) ds \right] dx \\ &\leq C \|u_n\|_{\mathcal{H}}^2 \left[\frac{(t - t_n)^2}{\Delta t} + (t - t_n) \right] \leq C \Delta t. \end{aligned}$$

This completes the proof. $\square$

7.1.2. Convergence of $u^{\Delta t}(t, x)$. Thanks to Lemma 7.2 and Lipschitz property of f and η , there exist u, f_u and η_u such that (up to a subsequence)

$$\begin{cases} u^{\Delta t} \rightharpoonup^* u & \text{in } L^\infty(0, T; L^2(\Omega \times \mathbb{R}^d)) \\ u^{\Delta t} \rightharpoonup u & \text{in } L^2((0, T) \times \Omega; H^1(\mathbb{R}^d)) \\ f(u^{\Delta t}) \rightharpoonup f_u & \text{in } L^2((0, T) \times \Omega; H^1(\mathbb{R}^d)) \\ \eta(u^{\Delta t}(\cdot - \Delta t); \cdot) \rightharpoonup \eta_u & \text{in } L^2(\Omega \times \Pi_T \times \mathbf{E}). \end{cases} \quad (\text{for fixed } \varepsilon > 0) \quad (7.5)$$

Let $v^{\Delta t}(t) = \sum_{k=1}^N \tilde{v}(t_k, \cdot) \mathbf{1}_{[t_{k-1}, t_k)}(t)$. Then, for any $\theta \in H^1(\mathbb{R}^d)$, we can rewrite (7.2), in terms of $u^{\Delta t}, \tilde{u}^{\Delta t}, \tilde{B}^{\Delta t}$ and $v^{\Delta t}$ as

$$\int_{\mathbb{R}^d} \left(\frac{\partial}{\partial t} (\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t})(t) \theta + \{ \varepsilon \nabla u^{\Delta t}(t) - v^{\Delta t}(t) f(u^{\Delta t}(t)) \} \cdot \nabla \theta \right) dx = 0. \quad (7.6)$$

In view of (7.6), one needs to show the boundedness of $\frac{\partial}{\partial t} (\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t})$ in $L^2(\Omega \times (0, T); H^{-1}(\mathbb{R}^d))$ and then identify the weak limit. Regarding this, we have the following lemma.

Lemma 7.4. *The sequence $\left\{ \frac{\partial}{\partial t} (\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t})(t) \right\}$ is bounded in $L^2(\Omega \times (0, T); H^{-1}(\mathbb{R}^d))$, and*

$$\frac{\partial}{\partial t} (\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t}) \rightharpoonup \frac{\partial}{\partial t} \left(u - \int_0^\cdot \int_{\mathbf{E}} \eta_u \tilde{N}(dz, ds) \right) \quad \text{in } L^2(\Omega \times (0, T); H^{-1}(\mathbb{R}^d))$$

where u is given by (7.5).

Proof. To prove the lemma, let $\Gamma = \Omega \times [0, T] \times \mathbf{E}$, $\mathcal{G} = \mathcal{P}_T \times \mathcal{L}(\mathbf{E})$ and $\varsigma = P \otimes \ell_t \otimes m$, where $\mathcal{P}_T$ represents predictable σ -algebra on $\Omega \times [0, T]$ and $\mathcal{L}(\mathbf{E})$ represents a Lebesgue σ -algebra on $\mathbf{E}$. Then $L^2((\Gamma, \mathcal{G}, \varsigma); \mathbb{R})$ consists of all square integrable predictable processes which are Borel measurable functions of z -variable. The space $L^2((\Gamma, \mathcal{G}, \varsigma); \mathbb{R})$ represents the space of square integrable predictable integrands for Itô-Lévy

integrals with respect to the compensated compound Poisson random measure $\tilde{N}(dz, dt)$. Moreover, Itô-Lévy integral defines a linear operator from $L^2((\Gamma, \mathcal{G}, \varsigma); \mathbb{R})$ to $L^2((\Omega, \mathcal{F}_T); \mathbb{R})$ and it preserves the norm (cf. for example [29]). Therefore, in view of (7.5) and the Lemma 7.3 along with above discussion, we have

$$\tilde{B}^{\Delta t} \rightharpoonup \int_0^\cdot \int_{\mathbf{E}} \eta_u \tilde{N}(dz, ds) \quad \text{in } L^2(\Omega \times \Pi_T).$$

Again, note that

$$\frac{\partial}{\partial t}(\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t})(t) = \sum_{k=1}^N \frac{(u_k - u_{k-1}) - (B_k - B_{k-1})}{\Delta t} \mathbf{1}_{[t_{k-1}, t_k]}.$$

From (7.2), we see that for any $\theta \in H^1(\mathbb{R}^d)$,

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\frac{u_{n+1} - u_n}{\Delta t} - \frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds) \right) \theta dx \\ &= -\varepsilon \int_{\mathbb{R}^d} \nabla u_{n+1} \cdot \nabla \theta dx - \int_{\mathbb{R}^d} \vec{v}(t_n, \cdot) f(u_{n+1}) \cdot \nabla \theta dx \\ &\leq \left\{ \varepsilon \|\nabla u_{n+1}\|_{L^2(\mathbb{R}^d)} + c_f V \|u_{n+1}\|_{L^2(\mathbb{R}^d)} \right\} \|\theta\|_{H^1(\mathbb{R}^d)}, \end{aligned}$$

and hence

$$\begin{aligned} & \sup_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\int_{\mathbb{R}^d} \left(\frac{u_{n+1} - u_n}{\Delta t} - \frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \int_{\mathbf{E}} \eta(u_n; z) \tilde{N}(dz, ds) \right) \theta dx}{\|\theta\|_{H^1(\mathbb{R}^d)}} \\ &\leq \varepsilon \|\nabla u_{n+1}\|_{L^2(\mathbb{R}^d)} + c_f V \|u_{n+1}\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

This implies that $\frac{\partial}{\partial t}(\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t})(t)$ is a bounded sequence in $L^2(\Omega \times (0, T); H^{-1}(\mathbb{R}^d))$.

To prove the second part of the lemma, we recall that $\tilde{B}^{\Delta t} \rightharpoonup \int_0^\cdot \int_{\mathbf{E}} \eta_u \tilde{N}(dz, ds)$ and $\tilde{u}^{\Delta t} \rightharpoonup u$ in $L^2(\Omega \times \Pi_T)$. In view of the first part of this lemma, one can conclude that, up to a subsequence

$$\frac{\partial}{\partial t}(\tilde{u}^{\Delta t} - \tilde{B}^{\Delta t}) \rightharpoonup \frac{\partial}{\partial t} \left(u - \int_0^\cdot \int_{\mathbf{E}} \eta_u \tilde{N}(dz, ds) \right) \quad \text{in } L^2(\Omega \times (0, T); H^{-1}(\mathbb{R}^d)).$$

This completes the proof. $\square$

In view of (7.5) and Lemma 7.4, one can pass to the limit in (7.6) and has, for $\theta \in H^1(\mathbb{R}^d)$

$$\int_{\mathbb{R}^d} \left(\frac{\partial}{\partial t} \left(u - \int_0^\cdot \int_{\mathbf{E}} \eta_u \tilde{N}(dz, ds) \right) \theta + \{ \varepsilon \nabla u(t) - \vec{v}(t, \cdot) f_u \} \cdot \nabla \theta \right) dx = 0.$$

An application of Itô-Lévy formula [11, Theorem A.1] to the function $e^{-ct} \|u(t)\|_2^2$ yields

$$\begin{aligned} & e^{-ct} \mathbb{E} [\|u(t)\|_2^2] + 2\varepsilon \int_0^t e^{-cs} \mathbb{E} [\|\nabla u(s)\|_2^2] ds - 2 \int_0^t \mathbb{E} \left[\int_{\mathbb{R}^d} e^{-cs} \vec{v}(s, x) f_u \cdot \nabla u dx \right] ds \\ &= \mathbb{E} [\|u_0\|_2^2] - c \int_0^t e^{-cs} \mathbb{E} [\|u(s)\|_2^2] ds + \mathbb{E} \left[\int_{\mathbf{E}} \int_0^t e^{-cs} \|\eta_u\|_2^2 ds m(dz) \right]. \end{aligned} \quad (7.7)$$

By choosing $\alpha > 0$ suitably in (7.3) and multiplying by e^{-ct_n} for positive $c > 0$, we have

$$\begin{aligned} & \mathbb{E} \left[\int_{\mathbb{R}^d} \left(e^{-ct_n} |u_{n+1}|^2 - e^{-ct_{n-1}} |u_n|^2 \right) dx \right] + 2\varepsilon \Delta t e^{-ct_n} \mathbb{E} \left[\int_{\mathbb{R}^d} |\nabla u_{n+1}|^2 dx \right] \\ &\leq \Delta t e^{-ct_n} \mathbb{E} \left[\int_{\mathbf{E}} \int_{\mathbb{R}^d} \eta^2(u_n; z) dx m(dz) \right] + (e^{-ct_n} - e^{-ct_{n-1}}) \mathbb{E} \left[\int_{\mathbb{R}^d} |u_n|^2 dx \right]. \end{aligned} \quad (7.8)$$

A straightforward calculation reveal that

$$\int_0^{t_{k+1}} e^{-cs} \mathbb{E} [\|\nabla u^{\Delta t}\|_2^2] ds \leq \sum_{n=0}^k e^{-ct_n} \mathbb{E} [\|\nabla u_{n+1}\|_2^2]$$

$$-ce^{-c\Delta t} \int_0^{t_k} e^{-cs} \mathbb{E}[\|u^{\Delta t}\|_2^2] ds \leq -c\Delta t \sum_{n=1}^k e^{-ct_{n+1}} \mathbb{E}[\|u_n\|_2^2]$$

and

$$\Delta t e^{-ct_n} \sum_{n=1}^k \mathbb{E} \left[\int_{\mathbf{E}} \int_{\mathbb{R}^d} \eta^2(u_n; z) dx m(dz) \right] \leq \int_0^{t_k} e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \int_{\mathbb{R}^d} \eta^2(u^{\Delta t}; z) dx m(dz) \right] ds + C\Delta t \mathbb{E}[\|u_0\|_2^2].$$

Therefore, by summing over n from 0 to k in (7.8), we obtain, for $t \in [t_k, t_{k+1})$

$$\begin{aligned} & e^{-ct} \mathbb{E}[\|u^{\Delta t}(t)\|_2^2] + 2\varepsilon \int_0^t e^{-cs} \mathbb{E}[\|\nabla u^{\Delta t}\|_2^2] ds \\ & \leq \mathbb{E}[\|u_0\|_2^2] + C\Delta t \mathbb{E}[\|u_0\|_2^2] + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \int_{\mathbb{R}^d} \eta^2(u^{\Delta t}; z) dx m(dz) \right] ds \\ & \quad - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E}[\|u^{\Delta t}\|_2^2] ds. \end{aligned} \quad (7.9)$$

Note that, for any $\theta \in H^1(\mathbb{R}^d)$ and any $s \in [0, T]$, there holds $\int_{\mathbb{R}^d} \vec{v}(s, x) f(\theta) \nabla \theta dx = 0$. Thus, using (7.9) we obtain

$$\begin{aligned} & e^{-ct} \mathbb{E}[\|u^{\Delta t}(t)\|_2^2] + 2\varepsilon \int_0^t e^{-cs} \mathbb{E}[\|\nabla(u^{\Delta t} - u)\|_2^2] ds \\ & \quad - 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} v(x, s) [f(u^{\Delta t}) - f(u)] \nabla(u^{\Delta t} - u) dx \right] ds \\ & \leq \mathbb{E}[\|u_0\|_2^2] + C\Delta t \mathbb{E}[\|u_0\|_2^2] + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \|\eta(u^{\Delta t}; z) - \eta(u; z)\|^2 m(dz) \right] ds \\ & \quad - \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \|\eta(u; z)\|^2 m(dz) \right] ds + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \int_{\mathbb{R}^d} \eta(u^{\Delta t}; z) \eta(u; z) dx m(dz) \right] ds \\ & \quad - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E}[\|u^{\Delta t} - u\|_2^2] ds + ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E}[\|u\|_2^2] ds + 2\varepsilon \int_0^t e^{-cs} \mathbb{E}[\|\nabla u\|_2^2] ds \\ & \quad - 2ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} u^{\Delta t} u dx \right] ds + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} \vec{v}(s, x) f(u^{\Delta t}) \nabla u dx \right] ds \\ & \quad + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} \vec{v}(s, x) f(u) \nabla u^{\Delta t} dx \right] ds - 4\varepsilon \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} \nabla u^{\Delta t} \nabla u dx \right] ds. \end{aligned} \quad (7.10)$$

In view of Young's inequality, one has

$$\begin{aligned} & -2\varepsilon \int_0^t e^{-cs} \mathbb{E}[\|\nabla(u^{\Delta t} - u)\|_2^2] ds + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbb{R}^d} \vec{v}(s, x) [f(u^{\Delta t}) - f(u)] \nabla(u^{\Delta t} - u) dx \right] ds \\ & \leq -\varepsilon \int_0^t e^{-cs} \mathbb{E}[\|\nabla(u^{\Delta t} - u)\|_2^2] ds + \frac{1}{\varepsilon} \int_0^t e^{-cs} \mathbb{E}[\|\vec{v}(s, \cdot) [f(u^{\Delta t}) - f(u)]\|_2^2] ds, \end{aligned} \quad (7.11)$$

and by choosing $c > 0$ with $\frac{1}{\varepsilon} V^2 C_f^2 + c_\eta \leq ce^{-c\Delta t}$, one arrive at

$$\begin{aligned} & \frac{1}{\varepsilon} \int_0^t e^{-cs} \mathbb{E}[\|\vec{v}(s, \cdot) [f(u^{\Delta t}) - f(u)]\|_2^2] ds + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \|\eta(u^{\Delta t}; z) - \eta(u; z)\|_2^2 m(dz) \right] ds \\ & \quad - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E}[\|u^{\Delta t} - u\|_2^2] ds \leq 0. \end{aligned} \quad (7.12)$$

We use (7.11)-(7.12) in (7.10) for the above choice of $c > 0$ along with (7.5) and (7.7) to have

$$\begin{aligned} & \limsup_{\Delta t} \int_0^T e^{-ct} \mathbb{E}[\|u^{\Delta t}(t)\|_2^2] dt + \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\mathbf{E}} \|\eta_u - \eta(u; z)\|_2^2 m(dz) \right] ds dt \\ & \leq \int_0^T e^{-ct} \mathbb{E}[\|u(t)\|_2^2] dt. \end{aligned}$$

Thus, we obtain $\eta_u = \eta(u; z)$ and $u^{\Delta t} \rightarrow u$ in $L^2(\Omega \times \Pi_T)$. Moreover, one can show that $f_u = f(u)$. Thus u is a weak solution to the viscous problem (7.1). Since it depends on $\varepsilon > 0$, we denote it by u_ε .

7.1.3. A-priori bounds for viscous solutions. Note that for fixed $\varepsilon > 0$, there exists a weak solution $u_\varepsilon \in H^1(\mathbb{R}^d)$ satisfying: $\mathbb{P}$ -a.s., and a.e. $t \in (0, T)$

$$\left\langle \frac{\partial}{\partial t} [u_\varepsilon - \int_0^t \int_{\mathbf{E}} \eta(u_\varepsilon(s, \cdot); z) \tilde{N}(dz, ds)], v \right\rangle + \int_{\mathbb{R}^d} \left\{ \vec{v}(t, x) f(u_\varepsilon(t, x)) + \varepsilon \nabla u_\varepsilon(t, x) \right\} \cdot \nabla v(x) dx = 0,$$

for any $v \in H^1(\mathbb{R}^d)$. We apply Itô-Lévy formula to $\beta(u) = u^2$, and then take expectation. The result is

$$\mathbb{E} \left[\|u_\varepsilon(t)\|_2^2 \right] + 2\varepsilon \int_0^t \mathbb{E} \left[\|\nabla u_\varepsilon\|_2^2 \right] ds \leq \mathbb{E} \left[\|u_\varepsilon(0)\|_2^2 \right] + C \int_0^t \mathbb{E} \left[\|u_\varepsilon(s)\|_2^2 \right] ds.$$

An application of Gronwall's inequality yields

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\|u_\varepsilon(t)\|_2^2 \right] + \varepsilon \int_0^T \mathbb{E} \left[\|\nabla u_\varepsilon(s)\|_2^2 \right] ds \leq C.$$

It is obvious from our existence proof that $\frac{\partial}{\partial t} \left[u_\varepsilon - \int_0^t \int_{\mathbf{E}} \eta(u_\varepsilon; z) \tilde{N}(dz, ds) \right] \in L^2(\Omega \times \Pi_T)$ and hence $\Delta u_\varepsilon \in L^2(\Omega \times \Pi_T)$. Moreover, if $u_0^\varepsilon \in L^{2p}(\mathbb{R}^d)$, $p \geq 1$, then a straightforward argumentation as in the proof of [2, Proposition A.5] gives $u_\varepsilon \in L^\infty(0, T; L^{2p}(\Omega \times \mathbb{R}^d))$.

The achieved results can be summarized into the following theorem.

Theorem 7.5. *Let $\varepsilon > 0$ is fixed and $u_0^\varepsilon \in H^1(\mathbb{R}^d)$. Then there exists a weak solution u_ε of (7.1) such that $\frac{\partial}{\partial t} \left[u_\varepsilon - \int_0^t \int_{\mathbf{E}} \eta(u_\varepsilon; z) \tilde{N}(dz, ds) \right], \Delta u_\varepsilon \in L^2(\Omega \times \Pi_T)$. Moreover the following estimate holds:*

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\|u_\varepsilon(t)\|_2^2 \right] + \varepsilon \int_0^T \mathbb{E} \left[\|\nabla u_\varepsilon(s)\|_2^2 \right] ds \leq C.$$

Furthermore, if $u_0^\varepsilon \in L^{2p}(\mathbb{R}^d)$, $p \geq 1$, then $u_\varepsilon \in L^\infty(0, T; L^{2p}(\Omega \times \mathbb{R}^d))$.

7.2. Proof of Theorem 2.1. In this subsection, we prove existence of generalized entropy solution in the sense of Definition 2.4. For this, fix a nonnegative test function $\psi \in C_c^\infty([0, \infty) \times \mathbb{R}^d)$, $B \in \mathcal{F}_T$ and convex entropy flux pair (β, ζ) . Now apply Itô-Lévy formula to (7.1) and conclude

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\mathbf{1}_B \int_{\mathbb{R}^d} \beta(u_0^\varepsilon(x)) \psi(0, x) dx \right] - \varepsilon \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \beta'(u_\varepsilon(t, x)) \nabla u_\varepsilon(t, x) \cdot \nabla \psi(t, x) dx dt \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \left(\beta(u_\varepsilon(t, x)) \partial_t \psi(t, x) + \nabla \psi(t, x) \cdot \vec{v}(t, x) \zeta(u_\varepsilon(t, x)) \right) dx dt \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \eta(u_\varepsilon(t, x); z) \beta'(u_\varepsilon(t, x) + \theta \eta(u_\varepsilon(t, x); z)) \psi(t, x) d\theta \tilde{N}(dz, dt) dx \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 (1 - \theta) \eta^2(u_\varepsilon(t, x); z) \beta''(u_\varepsilon(t, x) + \theta \eta(u_\varepsilon(t, x); z)) \psi(t, x) d\theta m(dz) dt dx \right] \end{aligned} \quad (7.13)$$

We use Young measure technique (cf. Subsection 6.2) to pass to the limit in (7.13) as $\varepsilon \rightarrow 0$. Moreover, there exists a $L^2(\mathbb{R}^d \times (0, 1))$ -valued predictable limit process $u \in L^\infty(0, T; L^2(\Omega \times \mathbb{R}^d \times (0, 1)))$ such that

$$\begin{aligned} &\mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_0^1 \left(\beta(u(t, x, \alpha)) \partial_t \psi(t, x) + \nabla \psi(t, x) \cdot \vec{v}(t, x) F^\beta(u(t, x, \alpha)) \right) d\alpha dx dt \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 \eta(u(t, x, \alpha); z) \beta'(u(t, x, \alpha) + \theta \eta(u(t, x, \alpha); z)) \psi(t, x) d\alpha d\theta \tilde{N}(dz, dt) dx \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \int_0^1 (1 - \theta) \eta^2(u(t, x, \alpha); z) \beta''(u(t, x, \alpha) + \theta \eta(u(t, x, \alpha); z)) \psi(t, x) d\alpha d\theta m(dz) dt dx \right] \\ &+ \mathbb{E} \left[\mathbf{1}_B \int_{\mathbb{R}^d} \beta(u_0(x)) \psi(0, x) dx \right] \geq 0. \end{aligned} \quad (7.14)$$

Since (7.14) holds for every $B \in \mathcal{F}_T$, we conclude that $\mathbb{P}$ -a.s., inequality (2.1) holds true as well. In other words, $u(t, x, \alpha)$ is a generalized entropy solution to the problem (1.1).

7.3. Proof of Theorem 2.2. To prove uniqueness of generalized entropy solutions, we follow the same arguments as in [9]. Let ρ and ϱ be the standard nonnegative mollifiers on $\mathbb{R}$ and $\mathbb{R}^d$ respectively such that $\text{supp}(\rho) \subset [-1, 0]$ and $\text{supp}(\varrho) = B_1(0)$. We define $\rho_{\delta_0}(r) = \frac{1}{\delta_0} \rho(\frac{r}{\delta_0})$ and $\varrho_\delta(x) = \frac{1}{\delta^d} \varrho(\frac{x}{\delta})$, where δ and δ_0 are two positive constants. Given a nonnegative test function $\psi \in C_c^{1,2}([0, \infty) \times \mathbb{R}^d)$ and two positive constants δ and δ_0 , we define

$$\phi_{\delta, \delta_0}(t, x, s, y) = \rho_{\delta_0}(t - s) \varrho_\delta(x - y) \psi(s, y).$$

Let $\beta : \mathbb{R} \rightarrow \mathbb{R}$ be a C^∞ function satisfying

$$\beta(0) = 0, \quad \beta(-r) = \beta(r), \quad \beta'(-r) = -\beta'(r), \quad \beta'' \geq 0,$$

and

$$\beta'(r) = \begin{cases} -1 & \text{when } r \leq -1, \\ \in [-1, 1] & \text{when } |r| < 1, \\ +1 & \text{when } r \geq 1. \end{cases}$$

For any $\vartheta > 0$, define $\beta_\vartheta : \mathbb{R} \rightarrow \mathbb{R}$ by $\beta_\vartheta(r) = \vartheta \beta(\frac{r}{\vartheta})$. Then

$$|r| - M_1 \vartheta \leq \beta_\vartheta(r) \leq |r| \quad \text{and} \quad |\beta_\vartheta''(r)| \leq \frac{M_2}{\vartheta} \mathbf{1}_{|r| \leq \vartheta},$$

where $M_1 = \sup_{|r| \leq 1} ||r| - \beta(r)|$ and $M_2 = \sup_{|r| \leq 1} |\beta''(r)|$. By simply dropping ϑ , for $\beta = \beta_\vartheta$ we define

$$\mathcal{F}^\beta(a, b) = \int_b^a \beta'(\sigma - b) f'(\sigma) d(\sigma).$$

Let $v(t, x, \alpha)$ be a generalized entropy solution of (1.1). Moreover, let ς be the standard symmetric nonnegative mollifier on $\mathbb{R}$ with support in $[-1, 1]$ and $\varsigma_l(r) = \frac{1}{l} \varsigma(\frac{r}{l})$ for $l > 0$. Given $k \in \mathbb{R}$, the function $\beta(\cdot - k)$ is a smooth convex function and $(\beta(\cdot - k), \mathcal{F}^\beta(\cdot, k))$ is a convex entropy pair.

Consider the entropy inequality for $v(t, x, \alpha)$, based on the entropy pair $(\beta(\cdot - k), \mathcal{F}^\beta(\cdot, k))$, and then multiply by $\varsigma_l(u_\varepsilon(s, y) - k)$, integrate with respect to s, y, k and take the expectation. The result is

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \beta(v_0(x) - k) \phi_{\delta, \delta_0}(0, x, s, y) \varsigma_l(u_\varepsilon(s, y) - k) dk dx dy ds \right] \\ &+ \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \beta(v(t, x, \alpha) - k) \partial_t \phi_{\delta, \delta_0}(t, x, s, y) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dt dy ds \right] \\ &+ \mathbb{E} \left[\int_{\Pi_T^2} \int_{\mathbb{R}} \int_{\mathbf{E}} \int_0^1 \left(\beta(v(t, x, \alpha) + \eta(v(t, x, \alpha); z) - k) - \beta(v(t, x, \alpha) - k) \right) \right. \\ &\quad \left. \times \phi_{\delta, \delta_0} dx d\alpha \tilde{N}(dz, dt) \varsigma_l(u_\varepsilon(s, y) - k) dk dy ds \right] \\ &+ \mathbb{E} \left[\int_{\Pi_T^2} \int_{\mathbf{E}} \int_{\mathbb{R}} \int_0^1 \left(\beta(v(t, x, \alpha) + \eta(v(t, x, \alpha); z) - k) - \beta(v(t, x, \alpha) - k) \right. \right. \\ &\quad \left. \left. - \eta(v(t, x, \alpha); z) \beta'(v(t, x, \alpha) - k) \right) \phi_{\delta, \delta_0} \varsigma_l(u_\varepsilon(s, y) - k) d\alpha dk dx m(dz) dt dy ds \right] \\ &+ \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(v(t, x, \alpha), k) \vec{v}(t, x) \cdot \nabla_x \phi_{\delta, \delta_0}(t, x, s, y) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dt dy ds \right] \\ &=: I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned} \tag{7.15}$$

Since $u_\varepsilon(s, y)$ is a viscous solution to the problem (7.1), one has

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \beta(u_\varepsilon(0, y) - k) \phi_{\delta, \delta_0}(t, x, 0, y) \varsigma_l(v(t, x, \alpha) - k) dk d\alpha dx dy dt \right] \\ &+ \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \beta(u_\varepsilon(s, y) - k) \partial_s \phi_{\delta, \delta_0}(t, x, s, y) \varsigma_l(v(t, x, \alpha) - k) dk d\alpha dy ds dx dt \right] \end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \left[\int_{\Pi_T^2} \int_{\mathbf{E}} \int_{\mathbb{R}} \int_0^1 \left(\beta(u_\varepsilon(s, y) + \eta_\varepsilon(y, u_\varepsilon(s, y); z) - k) - \beta(u_\varepsilon(s, y) - k) \right) \right. \\
& \quad \left. \times \phi_{\delta, \delta_0}(t, x, s, y) \varsigma_l(v(t, x, \alpha) - k) dy d\alpha dk \tilde{N}(dz, ds) dx dt \right] \\
& + \mathbb{E} \left[\int_{\Pi_T^2} \int_{\mathbf{E}} \int_{\mathbb{R}} \int_0^1 \left(\beta(u_\varepsilon(s, y) + \eta_\varepsilon(y, u_\varepsilon(s, y); z) - k) - \eta_\varepsilon(y, u_\varepsilon(s, y); z) \beta'(u_\varepsilon(s, y) - k) \right. \right. \\
& \quad \left. \left. - \beta(u_\varepsilon(s, y) - k) \right) \phi_{\delta, \delta_0}(t, x; s, y) \varsigma_l(v(t, x, \alpha) - k) d\alpha dk dy m(dz) ds dx dt \right] \\
& + \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(u_\varepsilon(s, y), k) \vec{v}(s, y) \cdot \nabla_y \varrho_\delta(x - y) \psi(s, y) \rho_{\delta_0}(t - s) \varsigma_l(v(t, x, \alpha) - k) dk d\alpha dx dt dy ds \right] \\
& + \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(u_\varepsilon(s, y), k) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) \rho_{\delta_0}(t - s) \varsigma_l(v(t, x, \alpha) - k) dk d\alpha dx dt dy ds \right] \\
& - \varepsilon \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} \beta'(u_\varepsilon(s, y) - k) \nabla_y u_\varepsilon(s, y) \cdot \nabla_y \phi_{\delta, \delta_0}(t, x, s, y) \varsigma_l(v(t, x, \alpha) - k) dk d\alpha dy ds dx dt \right] \\
& =: J_1 + J_2 + J_3 + J_4 + J_5 + J_6 + J_7. \tag{7.16}
\end{aligned}$$

We now add (7.15) and (7.16), and compute limits with respect to the various parameters involved. In [9], convergence of the terms $I_i (i = 1, 2, 3, 4)$ and $J_j (j = 1, 2, 3, 4, 7)$ has been studied in details. Therefore, we only study the terms involving flux function namely the terms I_5 , J_5 and J_6 in details.

We first consider the term $I_5 + J_5$ and prove the following lemma.

Lemma 7.6. *There holds*

$$\lim_{\delta \downarrow 0, \vartheta \downarrow 0, l \downarrow 0} \left[\lim_{\delta_0 \downarrow 0} (I_5 + J_5) \right] = 0.$$

Proof. Note that

$$\begin{aligned}
& \left| I_5 - \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(v(s, x, \alpha), k) \vec{v}(s, x) \cdot \nabla_x \varrho_\delta(x - y) \psi(s, y) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy ds \right] \right| \\
& \leq \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} |\mathcal{F}^\beta(v(t, x, \alpha), k) - \mathcal{F}^\beta(v(s, x, \alpha), k)| |\vec{v}(t, x)| |\nabla_x \varrho_\delta(x - y)| \psi(s, y) \right. \\
& \quad \left. \times \rho_{\delta_0}(t - s) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy dt ds \right] \\
& + \left| \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(v(s, x, \alpha), k) \vec{v}(s, x) \cdot \nabla_x \varrho_\delta(x - y) \psi(s, y) \left(1 - \int_{t=0}^T \rho_{\delta_0}(t - s) dt \right) \right. \right. \\
& \quad \left. \left. \times \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy ds \right] \right| \\
& + \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 \int_{\mathbb{R}} |\mathcal{F}^\beta(v(s, x, \alpha), k)| |\vec{v}(s, x) - \vec{v}(t, x)| |\nabla_x \varrho_\delta(x - y)| \psi(s, y) \right. \\
& \quad \left. \times \rho_{\delta_0}(t - s) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy ds dt \right] \\
& \leq \mathbb{E} \left[\int_{s=\delta_0}^T \int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} |\mathcal{F}^\beta(v(t, x, \alpha), k) - \mathcal{F}^\beta(v(s, x, \alpha), k)| |\vec{v}(t, x)| |\nabla_x \varrho_\delta(x - y)| \psi(s, y) \right. \\
& \quad \left. \times \rho_{\delta_0}(t - s) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy dt ds \right] + o(\delta_0) \\
& + C \mathbb{E} \left[\int_{s=0}^{\delta_0} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} |\mathcal{F}^\beta(v(s, x, \alpha), k) \vec{v}(s, x) \cdot \nabla_x \varrho_\delta(x - y)| \psi(s, y) \varsigma_l(u_\varepsilon(s, y) - k) dk d\alpha dx dy ds \right] \\
& \quad \text{(we have used the fact that } \int_0^T \rho_{\delta_0}(t - s) dt \leq 1, \text{ equality holds if } s \geq \delta_0) \\
& \leq C \mathbb{E} \left[\int_{s=\delta_0}^T \int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 |\nabla_x \varrho_\delta(x - y)| |v(t, x, \alpha) - v(s, x, \alpha)| \psi(s, y) \rho_{\delta_0}(t - s) d\alpha dx dy dt ds \right] + o(\delta_0)
\end{aligned}$$

(we have used the Lipschitz continuity of $\mathcal{F}^\beta(\cdot, k)$ in above)

$$\begin{aligned} &\leq C \left(\mathbb{E} \left[\int_{s=\delta_0}^T \int_{\Pi_T} \int_0^1 |v(t, x, \alpha) - v(s, x, \alpha)|^2 \rho_{\delta_0}(t-s) d\alpha dx dt ds \right] \right)^{\frac{1}{2}} + o(\delta_0) \\ &\leq C \left(\mathbb{E} \left[\int_{r=0}^1 \int_{\Pi_T} \int_0^1 |v(t + \delta_0 r, x, \alpha) - v(t, x, \alpha)|^2 \rho(-r) d\alpha dt dx dr \right] \right)^{\frac{1}{2}} + o(\delta_0). \end{aligned}$$

Note that, $\lim_{\delta_0 \downarrow 0} \int_{\Pi_T} \int_0^1 |v(t + \delta_0 r, x, \alpha) - v(t, x, \alpha)|^2 d\alpha dx dt \rightarrow 0$ almost surely for all $r \in [0, 1]$. Therefore, by the bounded convergence theorem,

$$\lim_{\delta_0 \downarrow 0} \mathbb{E} \left[\int_{r=0}^1 \int_{\Pi_T} \int_0^1 |v(t + \delta_0 r, x, \alpha) - v(t, x, \alpha)|^2 \rho(-r) d\alpha dx dr dt \right] = 0.$$

This implies that

$$\lim_{\delta_0 \rightarrow 0} I_5 = -\mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(v(s, x, \alpha), u_\varepsilon(s, y) - k) \vec{v}(s, x) \cdot \nabla_y \varrho_\delta(x - y) \psi(s, y) \varsigma_l(k) dk d\alpha dx dy ds \right].$$

In a similar manner, we find

$$\lim_{\delta_0 \rightarrow 0} J_5 = \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha) - k) \vec{v}(s, y) \cdot \nabla_y \varrho_\delta(x - y) \psi(s, y) \varsigma_l(k) d\alpha dk dx dy ds \right].$$

Note that, since $\operatorname{div}_x \vec{v}(t, x) = 0$ for all $(t, x) \in \Pi_T$, integration by parts formula yields

$$\begin{aligned} &\mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, y, \alpha) - k) [\vec{v}(s, x) - \vec{v}(s, y)] \cdot \nabla_y \varrho_\delta(x - y) \right. \\ &\quad \left. \times \psi(s, y) \varsigma_l(k) dk d\alpha dx dy ds \right] = 0. \end{aligned}$$

Hence, we have

$$\begin{aligned} \lim_{\delta_0 \rightarrow 0} (I_5 + J_5) &= \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \left(\mathcal{F}^\beta(v(s, x, \alpha), u_\varepsilon(s, y) - k) - \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha) - k) \right) \right. \\ &\quad \left. \times \vec{v}(s, x) \cdot \nabla_x \varrho_\delta(x - y) \psi(s, y) \varsigma_l(k) d\alpha dk dx ds dy \right] \\ &\quad + \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \left(\mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha) - k) - \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, y, \alpha) - k) \right) \right. \\ &\quad \left. \times (\vec{v}(s, x) - \vec{v}(s, y)) \cdot \nabla_x \varrho_\delta(x - y) \psi(s, y) \varsigma_l(k) d\alpha dk dx ds dy \right]. \end{aligned}$$

Define $\mathcal{F}(a, b) = \operatorname{sign}(a - b)(f(a) - f(b))$. Then, $\mathcal{F}$ is symmetric (i.e., $\mathcal{F}(a, b) = \mathcal{F}(b, a)$) and Lipschitz continuous in both of its variables. Moreover,

$$|\mathcal{F}^\beta(a, b) - \mathcal{F}(a, b)| \leq \vartheta c_f. \quad (7.17)$$

Therefore, one has

$$\left| \lim_{\delta_0 \rightarrow 0} (I_5 + J_5) \right| \leq C(c_f, v, \psi) \delta + C(c_f, V, \psi) \frac{\vartheta}{\delta} + C(c_f, V, \psi) \frac{l}{\delta}$$

and hence

$$\lim_{\delta \downarrow 0, \vartheta \downarrow 0, l \downarrow 0} \left[\lim_{\delta_0 \downarrow 0} (I_5 + J_5) \right] = 0.$$

□

Lemma 7.7. *It holds that*

$$\begin{aligned} J_6 &\xrightarrow{\delta_0 \rightarrow 0} \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(u_\varepsilon(s, y), k) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) \varsigma_l(v(s, x, \alpha) - k) dk d\alpha dx dy ds \right] \\ &\xrightarrow{l \rightarrow 0} \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\alpha dx dy ds \right] \end{aligned}$$

$$\begin{aligned}
& \xrightarrow{\varepsilon \rightarrow 0} \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_0^1 \mathcal{F}^\beta(u(s, y, \gamma), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\gamma d\alpha dx dy ds \right] \\
& \xrightarrow{\vartheta \rightarrow 0} \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_0^1 \mathcal{F}(u(s, y, \gamma), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\gamma d\alpha dx dy ds \right] \\
& \xrightarrow{\delta \rightarrow 0} \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \int_0^1 \mathcal{F}(u(s, y, \gamma), v(s, y, \alpha)) \cdot \nabla_y \psi(s, y) d\gamma d\alpha dy ds \right].
\end{aligned}$$

Proof. The proof is divided into five steps.

Step 1: We will justify the $\delta_0 \rightarrow 0$ limit. Define

$$\begin{aligned}
\mathcal{B}_1 &:= \left| J_6 - \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), k) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) \varsigma_l(v(s, x, \alpha) - k) d\alpha dk dy dx ds \right] \right| \\
&= \left| \mathbb{E} \left[\int_{\Pi_T^2} \int_{\mathbb{R}} \int_0^1 \left(\mathcal{F}^\beta(u_\varepsilon(s, y), v(t, x, \alpha) - k) - \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha) - k) \right) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \right. \right. \\
&\quad \left. \left. \times \rho_{\delta_0}(t - s) \varrho_\delta(x - y) \varsigma_l(k) d\alpha dk dy ds dx dt \right] \right. \\
&\quad \left. - \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha) - k) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) \right. \right. \\
&\quad \left. \left. \times \left(1 - \int_0^T \rho_{\delta_0}(t - s) dt \right) \varsigma_l(k) d\alpha dk dy dx ds \right] \right| \\
&\leq C \mathbb{E} \left[\int_{\Pi_T^2} \int_0^1 |\nabla_y \psi(s, y)| \rho_{\delta_0}(t - s) \varrho_\delta(x - y) |v(s, x, \alpha) - v(t, x, \alpha)| d\alpha dy ds dx dt \right] \\
&\quad + C \mathbb{E} \left[\int_0^{\delta_0} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \int_0^1 |\mathcal{F}^\beta(u_\varepsilon(s, y), k)| |\nabla_y \psi(s, y)| \varrho_\delta(x - y) \varsigma_l(v(s, x, \alpha) - k) d\alpha dk dy dx ds \right] \\
&\leq C \mathbb{E} \left[\int_{\delta_0}^T \int_{\mathbb{R}^d} \int_0^T \int_0^1 |v(s, x, \alpha) - v(t, x, \alpha)| \rho_{\delta_0}(t - s) d\alpha dt dx ds \right] + o(\delta_0) \\
&\leq C \left(\mathbb{E} \left[\int_{\delta_0}^T \int_0^T \int_{\mathbb{R}^d} \int_0^1 |v(s, x, \alpha) - v(t, x, \alpha)|^2 \rho_{\delta_0}(t - s) d\alpha dx dt ds \right] \right)^{\frac{1}{2}} + o(\delta_0) \longrightarrow 0 \text{ as } \delta_0 \rightarrow 0,
\end{aligned}$$

and therefore the first step follows.

Step 2: We will justify the $l \rightarrow 0$ limit. Let

$$\begin{aligned}
\mathcal{B}_2 &:= \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \mathcal{F}^\beta(u_\varepsilon(s, y), k) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) \varsigma_l(v(s, x, \alpha) - k) dk d\alpha dx dy ds \right] \\
&\quad - \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\alpha dx dy ds \right] \\
&= \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_{\mathbb{R}} \left(\mathcal{F}^\beta(u_\varepsilon(s, y), k) - \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha)) \right) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \right. \\
&\quad \left. \times \varrho_\delta(x - y) \varsigma_l(v(s, x, \alpha) - k) dk d\alpha dx dy ds \right].
\end{aligned}$$

By using boundedness of $\vec{v}$ and Lipschitz property of $\mathcal{F}^\beta$, we arrive at

$$|\mathcal{B}_2| \leq Cl \int_{\Pi_T} |\nabla_y \psi(s, y)| dy ds \rightarrow 0 \quad \text{as } l \rightarrow 0.$$

Step 3: We now justify the passage to the limit $\varepsilon \rightarrow 0$. Let

$$G_x(s, y, \omega, \xi) = \int_{\mathbb{R}^d} \int_0^1 \mathcal{F}^\beta(\xi, v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\alpha dx.$$

Then $G_x(s, y, \omega, \xi)$ is a Caratheodory function for every $x \in \mathbb{R}^d$ and $\{G_x(s, y, \omega, u_{\varepsilon_n}(s, y))\}_n$ is bounded and uniformly integrable. Thus, by Proposition 6.1 we conclude that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\int_{\mathbb{R}^d} \int_{\Pi_T} \int_0^1 \mathcal{F}^\beta(u_\varepsilon(s, y), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\alpha dx ds dy \right] \\ &= \mathbb{E} \left[\int_{\mathbb{R}^d} \int_{\Pi_T} \int_0^1 \int_0^1 \mathcal{F}^\beta(u(s, y, \gamma), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\gamma d\alpha dx ds dy \right]. \end{aligned}$$

Step 4: Justification of the limit $\vartheta \rightarrow 0$. Let

$$\begin{aligned} \mathcal{B}_3 := & \mathbb{E} \left[\int_{\mathbb{R}^d} \int_{\Pi_T} \int_0^1 \int_0^1 \left(\mathcal{F}^\beta(u(s, y, \gamma), v(s, x, \alpha)) - \mathcal{F}(u(s, y, \gamma), v(s, x, \alpha)) \right) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \right. \\ & \left. \times \varrho_\delta(x - y) d\gamma d\alpha dx ds dy \right]. \end{aligned}$$

In view of (7.17) and A.2, we see that $|\mathcal{B}_3| \leq C(V, c_f) \vartheta \int_{\Pi_T} |\nabla_y \psi(s, y)| dy ds \rightarrow 0$ as $\vartheta \rightarrow 0$.

Step 5: Justification of the limit $\delta \rightarrow 0$.

$$\begin{aligned} \mathcal{B}_4 := & \left| \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_0^1 \mathcal{F}(u(s, y, \gamma), v(s, x, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) \varrho_\delta(x - y) d\gamma d\alpha dx dy ds \right] \right. \\ & \left. - \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \int_0^1 \mathcal{F}(u(s, y, \gamma), v(s, y, \alpha)) \vec{v}(s, y) \cdot \nabla_y \psi(s, y) d\gamma d\alpha dy ds \right] \right| \\ & \leq C \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 \int_0^1 |\mathcal{F}(u(s, y, \gamma), v(s, x, \alpha)) - \mathcal{F}(u(s, y, \gamma), v(s, y, \alpha))| \right. \\ & \quad \left. \times |\nabla_y \psi(s, y)| \varrho_\delta(x - y) d\gamma d\alpha dx dy ds \right]. \end{aligned}$$

Since $\mathcal{F}$ is Lipschitz continuous in both of its variables, by Cauchy-Schwartz's inequality, we have

$$|\mathcal{B}_4| \leq C \left(E \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_0^1 |v(s, y, \gamma) - v(s, y + \delta z, \gamma)|^2 \varrho(z) d\gamma dy dz ds \right] \right)^{\frac{1}{2}} = o(\delta).$$

This completes the proof. $\square$

Following [9], we arrive at

Lemma 7.8. *The following hold:*

$$\begin{aligned} \lim_{(\delta, \vartheta, \varepsilon, l, \delta_0) \rightarrow 0} (I_1 + J_1) &= \mathbb{E} \left[\int_{\mathbb{R}^d} |v_0(x) - u_0(x)| \psi(0, x) dx \right]; \quad \limsup_{(\varepsilon, l, \delta_0) \rightarrow 0} |J_7| = 0. \\ \lim_{(\delta, \vartheta, \varepsilon, l, \delta_0) \rightarrow 0} (I_2 + J_2) &= \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \int_0^1 |u(s, y, \gamma) - v(s, y, \alpha)| \partial_s \psi(s, y) d\gamma d\alpha dy ds \right]. \\ \lim_{(l, \delta_0) \rightarrow 0} (I_4 + J_4) &= \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbf{E}} \int_0^1 \int_0^1 \left\{ \beta''(u_\varepsilon(s, y) - v(s, x, \alpha) + \lambda \eta(u_\varepsilon(s, y); z)) |\eta(u_\varepsilon(s, y); z)|^2 \right. \right. \\ & \quad \left. \left. + \beta''(v(s, x, \alpha) - u_\varepsilon(s, y) + \lambda \eta(v(s, x, \alpha); z)) |\eta(v(s, x, \alpha); z)|^2 \right\} \right. \\ & \quad \left. \times (1 - \lambda) \psi(s, y) \varrho_\delta(x - y) d\alpha d\lambda m(dz) dy dy ds \right]. \end{aligned}$$

Let us consider the stochastic integrals. Note that $J_3 = 0$. In view of Itô-Lévy formula, we see that

$$\begin{aligned} I_3 &= \mathbb{E} \left[\int_{\mathbb{R}} \int_{\Pi_T} J[\beta', \phi_{\delta, \delta_0}](s; y, k) \left(\int_{s-\delta_0}^s \varsigma_l(u_\varepsilon(\sigma, y) - k) \operatorname{div}(f(u_\varepsilon(\sigma, y)) \vec{v}(\sigma, y)) d\sigma \right) ds dy dk \right] \\ & \quad - \mathbb{E} \left[\int_{\mathbb{R}} \int_{\Pi_T} J[\beta', \phi_{\delta, \delta_0}](s; y, k) \left(\int_{s-\delta_0}^s \varsigma_l(u_\varepsilon(\sigma, y) - k) \varepsilon \Delta u_\varepsilon(\sigma, y) d\sigma \right) ds dy dk \right] \\ & \quad + \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}} \int_{r=s-\delta_0}^s \int_{\mathbb{R}^d} \int_{\mathbf{E}} \int_0^1 \left(\beta(v(r, x, \alpha) + \eta(v(r, x, \alpha); z) - k) - \beta(v(r, x, \alpha) - k) \right) \right. \end{aligned}$$

$$\begin{aligned}
& \times \left(\varsigma_l(u_\varepsilon(r, y) + \eta(u_\varepsilon(r, y); z) - k) - \varsigma_l(u_\varepsilon(r, y) - k) \right) \\
& \quad \times \rho_{\delta_0}(r - s) \psi(s, y) \varrho_\delta(x - y) d\alpha m(dz) dx dr dk dy ds \Big] \\
& + \mathbb{E} \left[\int_{\mathbb{R}} \int_{\Pi_T} J[\beta, \phi_{\delta, \delta_0}](s; y, k) \left\{ \int_{s-\delta_0}^s \int_{\mathbf{E}} \int_0^1 (1 - \lambda) |\eta(u_\varepsilon(\sigma, y); z)|^2 \right. \right. \\
& \quad \left. \left. \times \varsigma_l''(u_\varepsilon(\sigma, y) - k + \lambda \eta(u_\varepsilon(\sigma, y); z)) d\lambda m(dz) d\sigma \right\} dy ds dk \right] \\
& =: A_1^{l, \varepsilon}(\delta, \delta_0) + A_2^{l, \varepsilon}(\delta, \delta_0) + B^{\varepsilon, l} + A_3^{l, \varepsilon}(\delta, \delta_0),
\end{aligned}$$

where

$$\begin{aligned}
J[\beta, \phi_{\delta, \delta_0}](s; y, k) &:= \int_{\Pi_T} \int_{\mathbf{E}} \int_0^1 \left(\beta(v(r, x, \alpha) + \eta(v(r, x, \alpha); z) - k) - \beta(v(r, x, \alpha) - k) \right) \\
& \quad \times \phi_{\delta, \delta_0}(r, x, s, y) d\alpha \tilde{N}(dz, dr) dx.
\end{aligned}$$

Thanks to the assumption **A.2**, by following the arguments as in the proof of [9, Lemma 5.6], we infer that $A_1^{l, \varepsilon}(\delta, \delta_0)$, $A_2^{l, \varepsilon}(\delta, \delta_0)$, $A_3^{l, \varepsilon}(\delta, \delta_0) \rightarrow 0$ as $\delta_0 \rightarrow 0$, and

$$\begin{aligned}
\lim_{(l, \delta_0) \rightarrow 0} B^{l, \varepsilon}(\delta, \delta_0) &= \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \int_{\mathbf{E}} \int_0^1 \left\{ \beta(v(r, x, \alpha) + \eta(v(r, x, \alpha); z) - u_\varepsilon(r, y) - \eta(u_\varepsilon; z)) \right. \right. \\
& \quad - \beta(v(r, x, \alpha) + \eta(v(r, x, \alpha); z) - u_\varepsilon(r, y)) + \beta(v(r, x, \alpha) - u_\varepsilon(r, y)) \\
& \quad \left. \left. - \beta(v(r, x, \alpha) - u_\varepsilon(r, y) - \eta(u_\varepsilon(r, y); z)) \right\} \psi(r, y) \varrho_\delta(x - y) d\alpha m(dz) dx dy dr \right],
\end{aligned}$$

which yields

$$\begin{aligned}
& \lim_{l \rightarrow 0} \lim_{\delta_0 \rightarrow 0} \left((I_3 + J_3) + (I_4 + J_4) \right) \\
& = \mathbb{E} \left[\int_{\Pi_T} \int_{\mathbb{R}^d} \left(\int_{\mathbf{E}} \int_0^1 \int_0^1 b^2(1 - \theta) \beta''(a + \theta b) d\theta d\alpha m(dz) \right) \psi(t, y) \varrho_\delta(x - y) dx dy dt \right],
\end{aligned}$$

where $a = v(t, x, \alpha) - u_\varepsilon(t, y)$ and $b = \eta(v(t, x, \alpha); z) - \eta(u_\varepsilon(t, y); z)$. In view of **A.3**, one has (cf. proof of [9, Lemma 5.11]) $b^2 \beta''_\theta(a + \theta b) \leq 2(1 - \lambda^*)^{-2} \vartheta h_1^2(z)$, and thus

$$\limsup_{\delta \rightarrow 0, \vartheta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \left[\lim_{l \rightarrow 0} \lim_{\delta_0 \rightarrow 0} \left((I_3 + J_3) + (I_4 + J_4) \right) \right] = 0.$$

Finally, we add (7.15) and (7.16), and pass to the limits $\delta_0 \rightarrow 0$, $l \rightarrow 0$, $\varepsilon \rightarrow 0$, $\vartheta \rightarrow 0$ and $\delta \rightarrow 0$ to arrive at the following Kato inequality

$$\begin{aligned}
& \mathbb{E} \left[\int_{\mathbb{R}^d} |v_0(x) - u_0(x)| \psi(0, x) dx \right] + \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \int_0^1 |v(t, x, \alpha) - u(t, x, \gamma)| \partial_t \psi(t, x) d\alpha d\gamma dx dt \right] \\
& + \mathbb{E} \left[\int_{\Pi_T} \int_0^1 \int_0^1 \mathcal{F}(v(t, x, \alpha), u(t, x, \gamma)) \vec{v}(t, x) \cdot \nabla_x \psi(t, x) d\alpha d\gamma dx dt \right] \geq 0, \quad (7.18)
\end{aligned}$$

where $0 \leq \psi \in H^1([0, \infty) \times \mathbb{R}^d)$ with compact support. One can choose special test function ψ and $u_0 = v_0$ in (7.18) to conclude that $u(t, x, \gamma) = v(t, x, \alpha)$ for a.e. $(t, x) \in \Pi_T$ and a.e. $(\alpha, \gamma) \in (0, 1)^2$ (cf. proof of [9, Theorem 2.2]). This finishes the proof.

REFERENCES

- [1] E. J. Balder. Lectures on Young measure theory and its applications in economics. *Rend. Istit. Mat. Univ. Trieste*, 31 Suppl. 1:1-69, 2000.
- [2] C. Bauzet, G. Vallet, and P. Wittbold. The Cauchy problem for a conservation law with a multiplicative stochastic perturbation. *Journal of Hyperbolic Differential Equations*. 9 (2012), no 4, 661-709.
- [3] C. Bauzet. On a time-splitting method for a scalar conservation law with a multiplicative stochastic perturbation and numerical experiments. *J. Evol. Equ.* 14 (2014), no. 2, 333-356.
- [4] C. Bauzet, G. Vallet, and P. Wittbold. The Dirichlet problem for a conservation law with a multiplicative stochastic perturbation. *J. Funct. Anal.* 266 (2014), no. 4, 2503-2545.
- [5] C. Bauzet, G. Vallet and P. Wittbold. A degenerate parabolic-hyperbolic Cauchy problem with a stochastic force. *J. Hyperbolic Diff. Equ.* 12, (2015), no.3, 501-533.

- [6] C. Bauzet, J. Charrier, and T. Gallouët. Convergence of flux-splitting finite volume schemes for hyperbolic scalar conservation laws with a multiplicative stochastic perturbation. *Math. Comp.* 85 (2016), no. 302, 2777-2813.
- [7] C. Bauzet, J. Charrier, and T. Gallouët. Convergence of monotone finite volume schemes for hyperbolic scalar conservation laws with multiplicative noise. *Stoch. Partial Differ. Equ. Anal. Comput.* 4 (2016), no. 1, 150-223.
- [8] I. H. Biswas and A. K. Majee. Stochastic conservation laws: weak-in-time formulation and strong entropy condition. *Journal of Functional Analysis* 267 (2014), 2199-2252.
- [9] I. H. Biswas, K. H. Karlsen, and A. K. Majee. Conservation laws driven by Lévy white noise. *Journal of Hyperbolic Differential Equations*, no. 3, 581-654.
- [10] I. H. Biswas, U. Koley, and A. K. Majee. Continuous dependence estimate for conservation laws with Lévy noise. *J. Differential Equations* 259 (2015), no. 9, 4683-4706.
- [11] I. H. Biswas, A. K. Majee, and G. Vallet. On the Cauchy problem of a degenerate parabolic-hyperbolic PDE with Lévy noise. *submitted*. ArXiv: <http://arxiv.org/pdf/1604.05009.pdf>
- [12] G. Q. Chen, Q. Ding, and K. H. Karlsen. On nonlinear stochastic balance laws. *Arch. Ration. Mech. Anal.*, 204(3):707-743, 2012.
- [13] C. M. Dafermos. Hyperbolic conservation laws in continuum physics. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 325. Springer-Verlag, Berlin, 2000.
- [14] A. Debussche and J. Vovelle. Scalar conservation laws with stochastic forcing. *J. Funct. Analysis*, 259 (2010), 1014-1042.
- [15] A. Debussche and M. Hofmanová, and J. Vovelle. Degenerate parabolic stochastic partial differential equations: quasilinear case. *Annals of Prob.*, Vol. 44, No. 3, 1916-1955, 2016.
- [16] Z. Dong and T. G. Xu. One-dimensional stochastic Burgers equation driven by Lévy processes. *J. Funct. Anal.*, 243(2):631-678, 2007.
- [17] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. *Handbook of numerical analysis, Vol. VII, Handb. Numer. Anal.*, VII,713-1020. North-Holland, Amsterdam, 2000.
- [18] J. Feng and D. Nualart. Stochastic scalar conservation laws. *J. Funct. Anal.*, 255(2):313-373, 2008.
- [19] E. Godlewski and P.-A. Raviart. *Hyperbolic systems of conservation laws*, volume 3/4 of *Mathématiques & Applications (Paris) [Mathematics and Applications]*. Ellipses, Paris, 1991.
- [20] H. Holden and N. H. Risebro. Conservation laws with random source. *Appl. Math. Optim.*, 36(1997), 229-241.
- [21] J. U. Kim. On a stochastic scalar conservation law, *Indiana Univ. Math. J.* 52 (1) (2003) 227-256.
- [22] U. Koley, A. K. Majee and G. Vallet. Continuous dependence estimate for a degenerate parabolic-hyperbolic equation with Lévy noise. *Stoch. Partial Differ. Equ. Anal. Comput. (to appear)*.
- [23] U. Koley, A. K. Majee and G. Vallet. A finite difference scheme for conservation laws driven by Lévy noise. *submitted*. ArXiv: <http://arxiv.org/pdf/1604.07840.pdf>
- [24] K. H. Karlsen and E. B. Storrøsten. Analysis of a splitting method for stochastic balance laws. ArXiv: <http://arxiv.org/pdf/1601.02428v2.pdf>
- [25] I. Kröker and C. Rohde. Finite volume schemes for hyperbolic balance laws with multiplicative noise. *Applied Numerical Mathematics* 62, 441-456, 2012.
- [26] S. N. Kruzkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81(123): 228-255, 1970.
- [27] J. Málek, J. Nečas, M. Rokyta, and M. Růžička. *Weak and measure-valued solutions to evolutionary PDEs*, volume 13 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall, London, 1996.
- [28] E. Y. Panov. On measure-valued solutions of the Cauchy problem for a first-order quasilinear equations. *Investig. Mathematics*, 60(2):335-377, 1996.
- [29] S. Peszat and J. Zabczyk. *Stochastic partial differential equations with Lévy noise*, volume 113 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2007. An evolution equation approach.
- [30] G. Vallet. Stochastic perturbation of nonlinear degenerate parabolic problems. *Differential Integral Equations* 21 (2008), no. 11-12, 1055-1082.
- [31] A. I. Vol'pert. Generalized solutions of degenerate second-order quasilinear parabolic and elliptic equations. *Adv. Differential Equations*, 5(10-12):1493-1518, 2000.

(Ananta K. Majee)

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, D-72076 TÜBINGEN, GERMANY.

E-mail address: majee@na.uni-tuebingen.de